

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number  
**WO 01/16346 A1**

- (51) International Patent Classification<sup>7</sup>: C12P 7/22, (74) Agent: SEAY, Nicholas, J.; Quarles & Brady LLP, P.O. Box 2113, Madison, WI 53701-2113 (US).  
C12N 9/02, 9/14, 1/20, 15/00
- (21) International Application Number: PCT/US00/23878 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 30 August 2000 (30.08.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/151,440 30 August 1999 (30.08.1999) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 Walnut Street, 13th Floor, P.O. Box 7365, Madison, WI 53707-7365 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): SUTHERS, Patrick, F. [US/US]; 806 Olin Avenue, Apartment 1, Madison, WI 53715 (US). CAMERON, Douglas, C. [US/US]; 3590 Rainier Lane, North Plymouth, MN 55447 (US).
- Published:  
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/16346 A1

(54) Title: PRODUCTION OF 3-HYDROXYPROPIONIC ACID IN RECOMBINANT ORGANISMS

(57) Abstract: The production of 3-hydroxypropionic acid (3-HP) from glycerol in a bacterial host is described. 3-HP is a useful feedstock for the production of polymeric materials. The genetic engineering of a bacterial host with two enzymes is sufficient to enable production of 3-HP. One enzyme is a glycerol dehydratase and the other is an aldehyde dehydrogenase.

## PRODUCTION OF 3-HYDROXYPROPIONIC ACID IN RECOMBINANT ORGANISMS

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S. Provisional Patent Application S.N.  
5 60/151,440 filed August 30, 1999.

### STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

The research project which gave rise to the invention described in this patent  
application was supported by EPA grant R824726-01. The United States Government  
10 may have certain rights in this invention.

### BACKGROUND OF THE INVENTION

The technology of genetic engineering allows the transfer of genetic traits  
between species and permits, in particular, the transfer of enzymes from one species to  
others. These techniques have first reached commercialization in connection with high-  
15 value added products such as pharmaceuticals. The techniques of genetic engineering  
are equally applicable and cost effective when applied to genes and enzymes which can  
be used to make basic chemical feedstocks.

A metabolic pathway of interest exists in the bacteria *Klebsiella pneumoniae*,  
which has the ability to biologically produce 3 - hydroxypropionaldehyde from glycerol.  
20 Native microorganisms have the ability to produce 1,3 - propanediol from glycerol as  
well. Commercial interests are exploring the production of 1,3 - propanediol from  
glycerol or glucose, in recombinant organisms which have been engineered to express  
the enzymes necessary for 1,3 - propanediol production from other organisms.

3 - hydroxypropionic acid CAS registry Number [503-66-2] (abbreviated as 3-  
25 HP) is a three carbon non-chiral organic molecule. The IUPAC nomenclature name for

this molecule is propionic acid 3 - hydroxy. It is also known as 3 - hydroxypropionate,  $\beta$  - hydroxypropionic acid,  $\beta$  - hydroxypropionate, 3 - hydroxypropionic acid, 3 - hydroxypropanoate, hydracrylic acid, ethylene lactic acid,  $\beta$  -lactic acid and 2 - deoxyglyceric acid. Applications of 3-HP include the manufacture of absorbable  
5 prosthetic devices and surgical sutures, incorporation into beta-lactams, production of acrylic acid, formation of trifluoromethylated alcohols or diols, polyhydroxyalkonates, and co-polymers with lactic acid. 3-HP for commercial use is now commonly produced by organic chemical syntheses. The 3-HP produced and sold by these methods is relatively expensive, and it would be cost prohibitive to use it for the production of  
10 monomers for polymer production. As discussed below, some organisms are known to produce 3-HP. However, there is not yet available a catalog of genes from these organisms and thus the ability to synthesize 3-HP using the enzymes natively responsible for the synthesis of that molecule in the native hosts which produce it does not now exist.

15 In addition to its commercial utility, 3-HP it is found in a number of biological processes, notably including many naturally occurring bio-polymers. Poly(3 - hydroxybutyrate) (PHB) is the most abundant member of the microbial polyesters which contain hydroxy monomers termed polyhydroxyalkonates (PHAs). PHB has utility as a biodegradable thermoplastic material and the material was first produced industrially in  
20 1982.

The majority of published research on PHA's that contain 3-HP has concentrated on two bacterial sources: *Ralstonia eutropha* ("*Alcaligenes eutrophus*") and *Pseudomonas oleovorans*. Both *Ralstonia eutropha* and *Pseudomonas oleovorans* are able to grow on a nitrogen free media containing 3 - hydroxy - propionic acid, 1,5 -  
25 pentanediol or 1,7 - heptanediol. When 3-HP is the major hydroxy-acid added to the growth media, poly(3 - hydroxybutyrate - co - 3 - hydroxypropionic acid) is formed containing 7 mol % 3 - hydroxypropionic acid. These cells also store 3 mol %, 3 - hydroxypropionic acid poly(3 - butyrate - co - 3 - hydroxypropionic acid).

Recombinant systems have been used to create PHAs. An *E. coli* strain  
30 engineered to express PHA synthase from either *Ralstonia eutropha* or *Zoogloea ramigera* produced poly(3 - hydroxypropionic acid) when feed 1,3 - propanediol.

Skraly, F. A. "Polyhydroxyalkanoates Produced by Recombinant *E. coli*." Poster at Engineering Foundation Conference: Metabolic Engineering II, 1998. An *E. coli* strain that expressed PHA synthase (MBX820), when provided with the genes encoding glycerol dehydratase and 1,3 - propanediol dehydratase from *K. pneumonia*, and 4 -  
5 hydroxybutyral- CoA transferase from *Clostridium kluyveri*, synthesized PHB from glucose.

Glycerol dehydratase, found in the bacterial pathway for the conversion of glycerol to 1,3 - propanediol, catalyzes the conversion of glycerol to 3 - hydroxypropionaldehyde and water. This enzyme has been found in a number of  
10 bacteria including strains of *Citrobacter*, *Klebsiella*, *Lactobacillus*, *Enterobacter* and *Clostridium*. In the 1,3 - propanediol pathway a second enzyme 1,3 - propanediol oxido-reductase (EC 1.1.202) reduces 3 - hydroxypropanaldehyde to 1,3 - propanediol in a NADH dependant reaction. The pathway for the conversion of glycerol to 1,3 - propanediol has been expressed in *E. coli*. Tong et al., Applied and  
15 Environmental Microbiology 57 (12) 3541-3546. The genes responsible for the production of 1,3 - propanediol were cloned from the *dha* regulon of *Klebsiella pneumoniae*. Glycerol is transported into the cell by the glycerol facilitator, and then converted into 3 - hydroxy - propionaldehyde by a coenzyme B<sub>12</sub>- dependent dehydratase. *E. coli* lacks a native *dha* regulon, consequently *E. coli* cannot grow  
20 anaerobically on glycerol without an exogenous electron acceptor such as nitrate or fumarate.

Aldehyde dehydrogenases are enzymes that catalyze the oxidation of aldehydes to carboxylic acids. The genes encoding non-specific aldehyde dehydrogenases have been identified in a wide variety of organisms e.g.; *ALDH2* from *Homo sapiens*, *ALD4*  
25 from *Saccharomyces cerevisiae*, and from *E. coli* both *aldA* and *aldB*, to name a few. These enzymes are classified by co-factor usage, most require either AND<sup>+</sup>, or NADP<sup>+</sup> and some will use either co-factor. The genes singled out for mention here are able to act on a number of different aldehydes and it likely that they may be able to oxidize 3 - hydroxy - propionaldehyde to 3 - hydroxypropionic acid.

## BRIEF SUMMARY OF THE INVENTION

The present invention is intended to permit the creation of a recombinant microbial host which is capable of synthesizing 3-HP from a starting material of glycerol or glucose. The glycerol or glucose is converted to 3 -

5 hydroxypropionaldehyde (abbreviated as 3-HPA) which is then converted to 3-HP.

This process requires the so-called *dhaB* gene from *Klebsiella pneumoniae* which encodes the enzyme glycerol dehydratase any one of four different aldehyde dehydrogenase genes to convert 3-HPA to 3-HP. The four aldehyde dehydrogenase genes used were *aldA* from the bacterium *E. coli*, *ALDH2* from humans, *ALD4* from the  
10 yeast *Saccharomyces cerevisiae*, and *aldB* from *E. coli*. The yeast gene appeared to give the best results.

It is an object of the present invention to provide a genetic construct which encodes glycerol dehydratase and aldehyde dehydrogenase enzymes necessary for the production of 3 - hydroxypropionic acid from glycerol.

15 It is also an object of the present invention to provide a method for the production of 3 - hydroxypropionic acid from glycerol.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiment thereof and from the claims.

20 BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

Not applicable.

## DETAILED DESCRIPTION OF THE INVENTION

It is disclosed here that it is possible to introduce into a bacterial host genes encoding two enzymes and thus confer upon that host the ability to produce 3-HP from  
25 glycerol. The two necessary enzymes are glycerol dehydratase and aldehyde dehydrogenase. It is here reported that the two enzymes are both necessary and sufficient to enable a strain of a suitable host, such as a competent *E. coli* strain, to make 3-HP from glycerol. An exemplary gene encoding a glycerol dehydratase is known, the *dhaB* gene from *Klebsiella pneumoniae*, sequenced and rendered convenient to use.  
30 Several exemplary aldehyde dehydrogenases are known, and their sequences are

presented here. From this information, it becomes practical to confer upon a bacterial host the ability to convert glycerol into 3-HP in a commercially reasonable manner.

It was not apparent before the completion of the work described here that these two diverse enzymes could be produced in a common host to produce the ability to  
5 make 3-HP. There are many known aldehyde dehydrogenase enzymes and genes, and the enzymes are known to have varying substrate specificities and efficiencies. There was not evidence, prior to the work described here, that the aldehyde dehydrogenase enzyme would work on the 3-hydroxypropionaldehyde (3-HPA) substrate to create 3-HP. Without that knowledge, there was no data from which to predict the effectiveness  
10 of the 3-HP production studies described below. An additional uncertainty arises from the fact that the intermediate aldehyde, 3-HPA, is toxic to many bacterial host and thus the survival of the host is dependent upon the relative rates of enzymatic production and conversion of the aldehyde intermediate to non-toxic 3-HP.

A difficulty in the realization of the production of 3-HP desired here is that  
15 ribosome binding sites from non-native hosts are often ineffectual and lead to poor protein production and that many non-native promoters are often poorly transcribed and a bar to high protein expression. However, the inventors also recognized that a non-native promoter that is known to be very active and is inducible by the addition of a small molecule unrelated to the pathway being expressed is often a very efficient way to  
20 express and regulate the levels of enzymes expressed in hosts such as *E. coli*. To achieve high levels of regulated gene expression plasmids were constructed which placed the expression of all exogenous genes necessary for the production of 3 - hydroxypropionic acid from glycerol under the regulation of the *trc* promoter. The *trc* promoter, is efficient, not native to *E. coli*, and inducible by the addition of IPTG.

25 The present specification describes a genetic construct for use in the production of 3 - hydroxypropionic acid from glycerol. The genetic construct includes exemplary DNA sequences coding for the expression of a glycerol dehydratase and a DNA sequence coding for aldehyde dehydrogenase. The set of exemplary sequences necessary for the expression of glycerol dehydratase is collectively referred to as  
30 "*dhaB*". The set of sequences necessary for the expression of aldehyde dehydrogenase includes any one of four different genes which proved efficacious. The individual

aldehyde dehydrogenase sequences referred to individually as *ALDH4*, *ALD2*, *aldA* and *aldB*.

### Producing 3 - hydroxypropionic acid in a foreign host

In the work described below, the enzymes necessary for the production of 3 - hydroxypropionic acid from glycerol in *E. coli* were expressed under the regulation of the *trc* promoter, a non-native promoter inducible by the addition of IPTG. The glycerol dehydratase was encoded by the *dhaB* gene from *Klebsiella pneumoniae*, the aldehyde dehydrogenases used was any one of four different genes (*ALDH2* from *Homo sapiens*, *ALD4* from *S. cerevisiae*, *aldB* from *E. coli* or *aldA* from *E. coli*). Expression of these genes coding for glycerol dehydratase and any one of the genes encoding an aldehyde dehydrogenases was sufficient to enable the construct to produce 3-HP when the fermentation media was supplemented with glycerol. In all of these constructs, the *dhaB* gene was downstream from the gene encoding the aldehyde dehydrogenase used, and expression of both genes was regulated by the *trc* promoter. This order, however, is not required and the order of the gens on a construct and the use of multiple constructs is possible.

In a minimal genetic construct made based on the data presented here, the only genetic elements present that would be necessary are the structural genes *dhaB* and an aldehyde dehydrogenase gene encoding a protein that efficiently catalyzes the oxidation of 3-hydroxypropionaldehyde to 3-hydroxypropionic acid, and non-native promoter sequences specifically selected to give the type of inducible control most appropriate for the context of the process in which the construct is to be used. Extraneous pieces of DNA, whether retained in the construct or added from other DNA sequences, would not necessarily be detrimental to effective 3-HP synthesis by the host organism, but would not be needed. Each sequence to be translated would necessarily be preceded by a ribosome binding site, functional in the selected host so that the messenger RNA(s) coding for the proteins of interest could be translated by ribosomes. Terminator sequences immediately downstream of each translated unit would also be necessary in some organisms, particularly in eukaryotes. The construct could be part of an autonomously replicating sequence, such as a plasmid or phage vector, or could be

integrated into the genome of the host.

The structural genes and appropriate promoter(s) could be isolated by the use of restriction enzymes, by the polymerase chain reaction (PCR), by chemical synthesis of the appropriate oligonucleotides, or by other methods apparent to those skilled in the art or molecular biology. The promoter(s) would be derived from genomic DNA of other organism or from artificial genetic constructs containing promoters. Appropriate promoter fragments would be ligated into the construct upstream of the structural genes in any one of several possible arrangements.

The aldehyde dehydrogenase expressed would have: high specific activity towards 3-hydroxypropionaldehyde; be very stable in the host it is expressed in; be readily over expressed in the selected host; not be inhibited by either the substrates necessary for the reaction or the products formed by the reaction; be fully active under the fermentation conditions most favorable for the production of 3 - hydroxypropionic acid and be able to use either  $\text{NAD}^+$  or  $\text{NADP}^+$ .

One possible arrangement is the true operon, where one promoter is used to direct transcription in one direction of all necessary Open Reading Frames (ORFs). The entire message is then contained in one messenger RNA. The advantages of the operon are that it is relatively easy to construct, since only one promoter is needed; that it is relatively simple to replace the promoter with another promoter if that would be desirable later; and that it assures that the two genes are under the same regulation. The main disadvantage of the operon scheme is that the levels of the expression of the two genes cannot be varied independently. If it is found that the genes, for optimal 3 - hydroxypropionic acid synthesis, should be expressed at different levels, the operon in most cases cannot be used to realize this.

Another possible arrangement is the multiple-promoter scheme. Two or more promoters, with the same or distinct regulatory behavior, could be used to direct transcription of the genes. For example, one promoter could be used to direct transcription of *dhaB* and one to direct transcription of the gene encoding the appropriate aldehyde dehydrogenases. Because the genes theoretically can be transcribed and translated separately, a great number of combinations of multiple promoters is possible. Additionally, it would be most desirable to prevent the promoters



from interfering with one another. This could be achieved either by placing two promoters into the construct such that they direct transcription in opposite directions, or by inserting transcriptional terminator sequences downstream of each separately transcribed unit. The main advantage of the multiple-promoter construct is that it permits independent regulation of as many distinct units as desired, which could be important. The disadvantages are that it would be more difficult to construct; more difficult to amend later; and more difficult to effectively regulate, since multiple changes in fermentation conditions would need to be introduced and might render the performance of the fermentation somewhat less predictable.

- 10 In any construct, the promoter sequence(s) used should be functional in the selected host organism and preferably provide sufficient transcription of the genes comprising the glycerol to 3 - hydroxypropionic acid pathway to enable the construct to be adequately active in that host. The promoter sequence(s) used would also effect regulation of transcription of the genes enabling the glycerol to 3-HP pathway to be
- 15 adequately active under the fermentation conditions employed for 3-HP production, and preferably they would be inducible, such that expression of the genes could be modulated by the inclusion in, or exclusion from, the fermentation of a certain agents or conditions.

- A plausible example of the use of such a construct follows: one promoter, which
- 20 induced by the addition of an inexpensive chemical (the inducer) to the medium, could control transcription of both the *dhaB* gene and the gene encoding the appropriate aldehyde dehydrogenase. The cells would be permitted to grow in the absence of the inducer until they accumulated to a predetermined level. The inducer would then be added to the fermentation and nutritional changes commensurate with the altered
- 25 metabolism would be made to the medium as well. The cells would then be permitted to utilize the substrate(s) provided for 3-HP production (and additional biomass production if desired). After the cells could no longer use substrate to produce 3-HP, the fermentation would be stopped and the 3-HP recovered.

#### Genetic Sequences

- 30 To express glycerol dehydratase and a suitable aldehyde dehydrogenase, the two

enzymes necessary for the production of 3 - hydroxypropionic acid from glycerol, it is required that the DNA sequences containing the glycerol dehydratase and aldehyde dehydrogenase coding sequences be combined with at least a promoter sequence (preferably a non-native promoter although some native promoter activity may be present). An exemplary method of construction is described in the example below. To ensure that the present specification is enabling, the full sequences of the coding regions of genes for these enzymes is presented here.

Sequences 1, 3, 5 and 7 present different native genomic sequences for genes encoding aldehyde dehydrogenases.

10 SEQ ID NO:1 contains the full native DNA sequence encoding the *ALD4* enzyme from *Saccharomyces cerevisiae*. The amino acid sequence of the protein is presented as SEQ ID NO:2.

SEQ ID NO:3 includes the DNA sequence for the human *ALDH2* gene, again including the full protein coding region. The amino acid sequence for this human alcohol dehydrogenase is presented in SEQ ID NO:4.

15 SEQ ID NO:5 and 7 respectively present the full coding sequences from the *E. coli* genes *aldA* and *aldB*, both of which encode alcohol dehydrogenases. The amino acid sequences for the proteins encoded by the genes are presented in SEQ ID NO: 6 and 8 respectively.

20 SEQ ID NO:9 contains the native genomic DNA sequence for the *dhaB* gene from the *dha* regulon of *Klebsiella pneumoniae*. The coding sequences for this complex regulon produces five polypeptides, which are presented as SEQ ID NOS:10 through 13, which together provide the activity of the glycerol dehydratase enzyme.

Each of these coding sequences can be used to make genetic constructs for the expression of the appropriate enzymes in a heterologous hosts. In making genetic constructs for expression of the genes in such hosts, it is contemplated that heterologous promoters will be joined to the coding sequences for the enzymes, but all that it required is that the promoters be effective for the hosts in which the genes are to be expressed. It is also contemplated and envisioned that significant variations in DNA sequence are possible from the native DNA coding sequences presented here. As is well known in the art, due to the degeneracy of the genetic code, many different DNA sequences can

encode the expression of the same protein. So, when this document uses language specifying a DNA sequence encoding a protein, it is intended to encompass any DNA sequence which can be used to express that protein even if different from the genomic sequences presented here. It is also contemplated that conservative changes in the amino acid sequences of the proteins specified here can be made without departing from the present invention. In particular, deletions, additions and substitutions of one or more amino acids in a protein sequence can almost always be made without changing protein functionality. When the name of a protein is used here, it is intended to be equally applicable to both such minor changes in amino acid sequence and to allelic variations in native protein sequence as occurs within the species named as well as other closely related species.

It is possible that many of the above DNA sequences could be truncated and still express a protein that has the same enzymatic properties. One skilled in the art of molecular biology would appreciate that minor deletions, additions and mutations may not change the attributes of the designated base pair sequences; many of the nucleotide of the designated base pair sequences are probably not essential for their unique function. To determine whether or not an altered sequence or sequences has sufficient homology with the designated base pairs to function identically, one would simply create the candidate mutation, deletion or alteration and create a gene construct including the altered sequence together with promoter and termination sequences. This gene construct could be tested as, described below, for the production of 3-HP from glycerol.

Certain DNA primers were used to isolate or clone the genomic DNA sequences used in the experiments described below. While the sequence information presented here is sufficient to enable the construction of expression plasmids incorporating the genes identified here, in order to redundantly enable the use of these genes, primers which may be used to isolate the genes from their native hosts are described below.

The primers aldA\_L (SEQ ID NO:14), and aldA\_R (SEQ ID NO:15), were used to amplify the 1513 bp *aldA* fragment from genomic *E. coli* DNA (strain MG1655, a gift from the Genetic Stock Center, New Haven, CT). The gel purified PCR fragment containing a DNA sequence coding for the expression of aldehyde dehydrogenase was

inserted into *NcoI-XhoI* site of pSE380 (Invitrogen, San Diego, CA) to give pPFS3. The resulting plasmid contained *aldA* under the control of the *trc* promoter. This construct allowed for high-level expression of the *aldA* gene from *E. coli* under regulation of the *trc* promoter. Unless indicated otherwise all molecular biology and plasmid

5 constructions were done in *E. coli* AG1 (Stratagene, La Jolla, CA).

The primers aldB\_L (SEQ ID NO:20) and aldB\_R (SEQ ID NO:21), were used to amplify the 1574 bp *aldB* fragment from genomic *E. coli* DNA (strain MG1655). The resulting PCR converted the TGA stop codon into a TAA stop codon. The gel-purified PCR fragment containing the DNA sequence sufficiently coding for the  
10 expression of aldehyde dehydrogenase was inserted into the *KpnI-SacI* site of pSE380 to give pPFS12.

The primers ALD4\_L (SEQ ID NO : 16), and ALD4\_R (SEQ ID NO : 17), were used to amplify the 1595 bp ALD4 fragment from *S. cerevisiae* DNA (strain YPH500). The gel-purified fragment containing a DNA sequence coding for the expression of  
15 aldehyde dehydrogenase was inserted into the *KpnI-SacI* site of pPFS3 to give pPFS8. The resulting plasmid contained mature *ALD4* under control of the *trc* promoter.

The primers ALDH2\_L (SEQ ID NO:18), and ALDH2\_R (SEQ ID NO:19), were used to amplify the 1541 bp ALDH2 fragment from pT7-7::ALDH2, a gift from H. Weiner (Purdue University, West Lafayette, IN). The gel purified PCR fragment  
20 containing a DNA sequence sufficiently homologous to base pairs 22 to 1524, inclusive of SEQ ID NO : 3 so as to code for the expression of aldehyde dehydrogenase was inserted in to the *KpnI-SacI* site of pSE380 to give pPFS7. This sequence was moved from pPFS7 into the *KpnI-SacI* site of pPFS3 to give pPFS9. The resulting plasmid contained mature ALDH2 under the control of the *trc* promoter.

25 The primers pTRC\_L (SEQ ID NO:22), and pTRC\_R (SEQ ID NO:23), were used to amplify the 540 bp fragment from pSE380. The gel purified PCR fragment was inserted into the *HpaI-KpnI* site of pPFS3 to give pPFS13. The resulting plasmid deleted the "native" ribosome binding site of pSE380 and a *NcoI* site (which contained an extraneous ATG start codon upstream of the cloned genes). The *KpnI-SacI*  
30 fragments of pPFS8, pPFS9, and pPFS12 were inserted into the *KpnI-SacI* site of pPFS13 to give pPFS14, pPFS15, and pPFS16, respectively.

Assay for production of 3-HP

The efficacy of changes made as contemplated herein can be checked by the following tests. To test for the production of 3-HP, fermentation products can be quantified with a Waters Alliance Integrity HPLC system (Milford, MA) equipped with a refractive index detector, a photodiode array detector, and an Aminex HPX-87H (Bio-Rad, Hercules, CA) organic acids column. The mobile phase should be 0.01 N sulfuric acid solution (pH 2.0) at a flow rate of 0.5 mL/min. The column temperature should be set to 40°C. Compounds can be identified by determining if they co-elute with authentic standards. Prior to analysis, all samples should be filtered through 0.45 µM pore size membrane. (Gelman Sciences, Ann Arbor, MI). The fractions of the fermentation products collected using HPLC should be analyzed on a Varian Star 3400 CX, gas - chromatograph coupled to a Varian Saturn 3 mass spectrometer (GC-MS) (Walnut Creek, CA).

Assay for enzyme activity.

Aldehyde dehydrogenase activity can be determined by measuring the reduction of  $\beta$ -NAD<sup>+</sup> at 25°C with 3 - hydroxypropionaldehyde as a substrate. All buffers should contain 1 mM ethylenediaminetetraacetic acid (EDTA), 0.1 mM Pefabloc SC (Boehringer Mannheim, Indianapolis, IN) and 1 mM Tris (carboxyethyl) phosphine hydrochloride (TCEP-HCL). For ALD4, the solution should contain 100 mM Tris HCL Buffer (pH 8.0), 100 mM KCl. For ALDH2 the solution should contained 100 mM sodium pyrophosphate (pH 9.0). For AldA and AldB, the solution should contain 20 mM sodium glycine (pH 9.5). A total of 3.0 mL of buffer should be added to quartz cuvettes and allowed to equilibrate to assay temperature. From 5 to 20 µL of cell extract should be added and background activity recorded after the addition of  $\beta$ -NAD<sup>+</sup> to a final concentration of 0.67 mM. The reaction should be started by the addition of substrate (either acetaldehyde, propionaldehyde, or 3 - hydroxypropionaldehyde) to a final concentration of 2 mM. Assay mixtures should be stirred with micro-stirrers during the assays.

For aldehyde dehydrogenase activity assays, one unit is defined as the reduction

of 1.0  $\mu\text{M}$  of  $\beta\text{-AND}^+$  per minute at 25° C. These reactions can be monitored by following the change in absorbance at 340 nm ( $A_{340}$ ) at 25°C on a Varian Carry-1 Bio spectrophotometer (Sugar Land, TX). Total protein concentrations in the cell extracts can be determined using the Bradford assay method (Bio-Rad, Hercules, CA) with  
5 bovine serum albumin as the standard.

## EXAMPLES

### Plasmid constructions.

*Klebsiella pneumoniae* expresses glycerol dehydratase, an enzyme that catalyzes the conversion of glycerol to 3 - hydroxypropionaldehyde, (*dhaB*) and 1,3 -  
10 propanediol oxidoreductase an enzyme that catalyzes the conversion of 3 - hydroxypropionaldehyde to 1,3 - propanediol respectively (the gene product from *dhaT*). A plasmid encoding these two genes was created and expressed in *E. coli* (plasmid pTC53). The *dhaT* gene was deleted from pTC53 to create pMH34. The resulting plasmid still contained the DNA sequence complementary to base pairs 330 to  
15 2153 inclusion of SEQ ID NO : 9, the complement of base pairs 2166 to 2591, inclusive, of SEQ ID NO : 9, and the complement of base pairs 3191 to 4858, inclusive, of SEQ ID NO : 9, so as to code for the expression of glycerol dehydratase. The fragment of DNA encoding these sequences was excised from pMH34 by cutting it with *Sall-XbaI*, and the resulting fragment was gel purified (the purified fragment was gift  
20 from M. Hoffman of the University of Wisconsin - Madison). This DNA fragment was inserted into the *Sall-XbaI* site of pPFS13 to give pPFS17.

The resulting plasmid contained both the *aldA* and *dhaB* genes under the control of the *trc* promoter. Similarly, the gel-purified *Sall-XbaI* fragment from pMH34 was inserted into the *Sall-XbaI* sites of pPFS14, pPFS15, and pPFS16 to give pPFS18,  
25 pPFS19, and pPFS20, respectively. These plasmids contained *ALD4*, *ALDH2*, and *aldB*, respectively, as well as *dhaB* under the control of the *trc* promoter; in all of the constructs the *dhaB* gene were downstream of the gene encoding the aldehyde dehydrogenase.

### Expression in *E. coli*.

The efficacy of *E. coli* as a platform for the production of 3-HP from growth on glucose has been examined using a mathematical model developed for this purpose. The model was executed in two different ways assuming the conversion of one mole of  
5 glucose under either anaerobic or aerobic conditions either directly to 3-HP or to the production of 3-HP and ATP. The optimum yield under anaerobic conditions is 1 mole of 3-HP and 1 mole of lactate. The more realistic yield under anaerobic conditions is 0.5 moles of 3-HP, 1.5 moles of lactate and 1 mole of ATP. The optimum yield under aerobic conditions is 1.9 moles of 3-HP and 0.3 moles of CO<sub>2</sub>. The realistic yield under  
10 aerobic conditions is 1.85 moles of 3-HP, 0.35 moles of CO<sub>2</sub> and 1 mole of ATP.

The effect of 3-HP concentration on *E. coli* strain MG1655 growth was measured. Cells were grown on standard media with and without the addition of up to 80g/L of 3-HP. The best fit of these data demonstrated that 3-HP was only 1.4 times as inhibitory as lactic acid on the growth of *E. coli*. It is possible to economically produce  
15 lactic acid using *E. coli*, since 3-HP is only 1.4 times more inhibitory than lactic acid, it should be possible to use *E. coli* as a host for the commercial production of 3-HP.

### Media and growth conditions

The standard media contained the following per liter: 6 g Na<sub>2</sub>HPO<sub>4</sub>, 3 g KH<sub>2</sub>PO<sub>4</sub>, 1 g NH<sub>4</sub>Cl, 0.5 g NaCl, 3 mg CaCl<sub>2</sub>, 5 g yeast extract (Difco Laboratories, Detroit, MI)  
20 and 2 mM MgSO<sub>4</sub>. When necessary to retain plasmids ampicillin (100 mg/mL) was added to the media. Isopropyl-β-thiogalactopyranoside (IPTG) was added in varying amounts to induce gene expression. All fermentations were carried out in an incubator-shaker at 37 C and 200 rpm. Anaerobic fermentations were carried out in 500-mL anaerobic flasks with 300 mL of working volume. Inocula for fermentations were  
25 grown overnight in Luria-Bertani medium supplemented with ampicillin is necessary. The 300-mL fermentations were inoculated with 1.5 mL of the overnight culture. For enzyme assays, fermentations were incubated for 24 hours.

### Over expression of aldehyde dehydrogenase in *E. coli*.

Cells were harvested by centrifugation at 3000 x g for 10 minutes at 4°C with a

Beckman (Fullerton, CA) model J2-21 centrifuge. Cell pellets were washed twice in 100 mM potassium phosphate buffer at pH 7.2 and re-suspended in appropriate assay resuspension buffer equal to 5 x of the volume of the wet cell mass. The cells were homogenized using a French pressure cell. The homogenate was centrifuged at 40000 x g for 30 minutes. The supernatant was dialyzed against the appropriate resuspension buffer using 10000 molecular weight cut-off pleated dialysis tubing (Pierce, Rockford, IL) at 4°C. Dialysis buffer was changed after 2 hours, and 4 hours, and dialysis was stopped after being allowed to proceed overnight.

*E. coli* AG1 cells transfected with the plasmids constructed to express the *aldA*, *ALD4*, *ALDH2*, or *aldB* genes were grown in 500-mL anaerobic flasks. Twelve hours after the fermentations were inoculated IPTG was added to induce enzyme expression. The cells were allowed to grow for an additional 12 hours then harvested and lysed as discussed above. The soluble fraction of the lysate was assayed for aldehyde dehydrogenase activity using the substrate 3-hydroxypropionaldehyde in the buffer appropriate for the particular enzyme expressed. The plasmid, aldehyde dehydrogenase expressed and specific activity measured (U/mg of protein) were as follows: pPFS13, *aldA*, 0.2; pPFS14, *ALD4*, 0.5, pPFS15, *ALDH2*, 0.3; and pPFS16, *aldB*, 0.1. The control, *E. coli* strain AG1 harboring plasmid pSE380, encoded no exogenous aldehyde dehydrogenase activity and it had no detectable activity with 3-HP as substrate. It is clear from the activity assays that all four aldehyde dehydrogenases were expressed in *E. coli*. The aldehyde dehydrogenase cloned from *Saccharomyces cerevisiae* (*ADH4*) had the highest activity when 3-hydroxypropionaldehyde was used as the substrate (0.5 units/mg of protein).

*E. coli* cells transformed with plasmids expressing: aldehyde dehydrogenase; both aldehyde dehydrogenase and glycerol dehydratase, or neither gene; were grown and assayed for their ability to produce 3-HP from glycerol. The cells were grown on standard media supplemented with 6 µM of Coenzyme B<sub>12</sub>, under anaerobic conditions in the absence of light (to protect the integrity of the Coenzyme B<sub>12</sub> necessary for DhaB activity). After 12 hours, IPTG was added to induce expression of the genes under the *trc* promoter at the same time 5g/L of glycerol was added. After 12 more hours of anaerobic fermentation the fermentation broth was assayed for 3 - HP by HPLC and GC,



the plasmid, aldehyde dehydrogenase gene expressed and g/L of 3- HP measured were as follows: pSF17, *aldA*, 0.031; pPSF18 *ALD4*, 0.173; and pPSF19, *ALDH2*, 0.061. Cells expressing *dhaB* but no exogenous aldehyde dehydrogenase genes (plasmid pMH34) produced 0.015 g/L of 3 - HP. Cells expressing *aldA*, *ALD4*, *ALDH2* or *aldB* but not *dhaB* (plasmids pPFS13, pPFS14, pPFS15, pPFS16, respectively) all produced less than 0.005 g/L of 3-HP when the media the cells were growing in was supplemented with 2.5g/L of 3-hydroxypropionaldehyde.

#### Other Hosts and Promoters

Applications of the 3 - hydroxypropionic acid pathway such as the genetic constructs of the present invention can easily be expressed in other organisms. The required genes would need to be placed under control of an appropriate promoter or promoters. Some organism such as yeasts may require transcription terminators to be placed after each transcribed unit. The knowledge of the present intention makes such amendments possible. Such a genetic construct would need to be part of a vector that could either replicate in the new host or integrate into the chromosome of the new host. Many such vectors are commercially available for expression in gram-negative and gram-positive bacteria, yeast, mammalian cells, insect cell, plant, etc. For example, to express the 3-hydroxypropionic acid pathway in *Rhodobacter capsulatus*, one could obtain vector pNH2 from the American Type Culture Collection ( ATTC). This is a shuttle vector for use in *R. capsulatus* and *E. coli*. Organisms such as *Saccharomyces cerevisiae* which can convert glucose to glycerol could be used as a host, such a construct would enable the production of 3 - HP directly from glucose. Additionally, other substrates such as xylan could also be used given the selection of an appropriate host.

Stoichiometric analysis shows that best stoichiometric yield of 3-HP production in *E. coli* calculated on the basis of glucose consumed is obtained under aerobic conditions. Under aerobic condition CO<sub>2</sub> is the only carbon-containing co-product, in particular the generation of lactic acid which occurs under anaerobic conditions is avoided. Production of 3-HP under these conditions could result in a more economical recovery of 3-HP from the fermentation broth.

Alternatively, the *dhaB* gene and a gene encoding the appropriate aldehyde dehydrogenase could be cloned into the multiple cloning site of this vector in *E. coli* to facilitate construction, and then transformed into *R. capsulatus*. The *R. capsulatus* *nifH* promoter, provided on the plasmid, could be used to direct the transcription in *R.*  
5 *capsulatus* of the genes placed into pNF2 in series with one promoter, or with two copies of the *nifH* promoter. Expression of the genes in other organisms would require a procedure analogous to that presented here.

#### Alternative Aldehyde Dehydrogenases and Glycerol Dehydratases

Applications of the pathway for the production of 3-hydroxypropionic acid from  
10 glycerol can be made using other suitable aldehyde dehydrogenases. To be functional in this pathway an aldehyde dehydrogenase needs to be stable, readily expressed in the host of choice and have high enough activity towards 3-hydroxypropionaldehyde to enable it to make 3-HP. The knowledge of the present invention makes such amendments possible. A program of directed evolution could be undertaken to select  
15 for suitable aldehyde dehydrogenases or they could be recovered from native sources, the genes encoding these enzymes in conjunction with a gene encoding an appropriate glycerol dehydratase activity, would then be made part of any of the constructs envisioned here to produce 3 - hydroxypropionic acid from glycerol.

A similar program of enzyme improvement including for example directed  
20 evolution could be carried out using the *dhaB* gene from *Klebsiella pneumoniae* as a starting point to obtain other variants of glycerol dehydratase that are superior in efficiency and stability to the form used in this invention. Alternatively, enzymes which catalyzes the same reaction may be isolated from others organisms and used in place of the *Klebsiella pneumoniae* glycerol dehydratase. Such enzymes may be especially  
25 useful in alternative hosts wherein they may be more readily expressed, be more stable and more efficient under the fermentation conditions best suited to the growth of the construct and the production and recovery of 3-HP.

## CLAIM OR CLAIMS

I/WE CLAIM:

1. A method for producing 3-hydroxypropionic acid comprising the steps of  
providing in a fermenter a recombinant microorganism which expresses genes  
5 for non-native enzymes which are capable of catalyzing the production of 3-  
hydroxypropionic acid from glycerol;  
providing a source of glycerol or glucose for the recombinant microorganism,  
and  
fermenting the microorganism under conditions which result in the accumulation  
10 of 3-hydroxypropionic acid.
  
2. A method for producing 3-hydroxypropionic acid comprising the steps of  
providing in a fermenter a recombinant microorganism which carries genetic  
constructions for the expression of a glycerol dehydratase and an aldehyde  
dehydrogenase which are capable of catalyzing the production of 3-hydroxypropionic  
15 acid from glycerol;  
providing a source of glycerol or glucose for the recombinant microorganism,  
and  
fermenting the microorganism under conditions which result in the accumulation  
of 3-hydroxypropionic acid.

3. A method for producing 3-hydroxypropionic acid comprising the steps of providing in a fermenter a recombinant microorganism which carries a genetic construct which expresses the *dhaB* gene from *Klebsiella pneumoniae* and a gene for an aldehyde dehydrogenase, which are capable of catalyzing the production of 3-hydroxypropionic acid from glycerol;
- 5 providing a source of glycerol or glucose for the recombinant microorganism, and fermenting the microorganism under conditions which result in the accumulation of 3-hydroxypropionic acid.
- 10 4. The method of claim 3 wherein the gene for the aldehyde dehydrogenase is selected from the group consisting of *ALDH4*, *ALD2*, *aldA* and *aldB*.
5. The method of claim 3 wherein the aldehyde dehydrogenase is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.
- 15 6. A recombinant *E. coli* host comprising in its inheritable genetic materials foreign genes encoding a glycerol dehydratase and an aldehyde dehydrogenase, such that the host is capable of producing 3-hydroxypropionic acid from glycerol.
7. A recombinant *E. coli* host comprising in its inheritable genetic materials the *dhaB* gene from *Klebsiella pneumoniae* and the *ald4* gene from *Saccharomyces cerevisiae*, such that the host is capable of producing 3-hydroxypropionic from glycerol.

8. A bacterial host comprising in its inheritable genetic material a genetic construction encoding for the expression of a glycerol dehydratase enzyme and an aldehyde dehydrogenase enzyme, such that the bacterial host is capable of converting glycerol to 3-hydroxypropionic acid.

5           9. The bacterial host of claim 8 wherein the glycerol dehydratase from *Klebsiella pneumoniae*.

10           10. The bacterial host of claim 8 wherein the gene encoding the glycerol dehydratase is the *dhaB* gene from *Klebsiella pneumoniae*.

11. The bacterial host of claim 8 wherein the aldehyde dehydrogenase is  
10 selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

12. The bacterial host of claim 8 wherein the gene for the aldehyde dehydrogenase is selected from the group consisting of *ALDH4*, *ALD2*, *aldA* and *aldB*.

## SEQUENCE LISTING

<110> Suthers, Patrick F  
Cameron, Douglas C.

<120> Production of 3-Hydroxypropionic Acid in Recombinant Organisms

5 <130> UW960296.96617

<140>

<141>

<160> 23

<170> PatentIn Ver. 2.1

10 <210> 1

<211> 1529

<212> DNA

<213> Saccharomyces cerevisiae

<220>

15 <221> CDS

<222> (25)..(1509)

<400> 1

gtcgcggtac caaggaggta tcat atg tca cac ctt cct atg aca gtg cct 51

Met Ser His Leu Pro Met Thr Val Pro

20

1

5

atc aag ctg ccc aat ggg ttg gaa tat gag caa cca acg ggg ttg ttc 99

Ile Lys Leu Pro Asn Gly Leu Glu Tyr Glu Gln Pro Thr Gly Leu Phe

10

15

20

25

atc aac aac aag ttt gtt cct tct aaa cag aac aag acc ttc gaa gtc 147

25 Ile Asn Asn Lys Phe Val Pro Ser Lys Gln Asn Lys Thr Phe Glu Val

30

35

40

att aac cct tcc acg gaa gaa gaa ata tgt cat att tat gaa ggt aga 195  
 Ile Asn Pro Ser Thr Glu Glu Glu Ile Cys His Ile Tyr Glu Gly Arg  
 45 50 55

gag gac gat gtg gaa gag gcc gtg cag gcc gcc gac cgt gcc ttc tct 243  
 5 Glu Asp Asp Val Glu Glu Ala Val Gln Ala Ala Asp Arg Ala Phe Ser  
 60 65 70

aat ggg tct tgg aac ggt atc gac cct att gac agg ggt aag gct ttg 291  
 Asn Gly Ser Trp Asn Gly Ile Asp Pro Ile Asp Arg Gly Lys Ala Leu  
 75 80 85

10 tac agg tta gcc gaa tta att gaa cag gac aag gat gtc att gct tcc 339  
 Tyr Arg Leu Ala Glu Leu Ile Glu Gln Asp Lys Asp Val Ile Ala Ser  
 90 95 100 105

atc gag act ttg gat aac ggt aaa gct atc tct tcc tcg aga gga gat 387  
 Ile Glu Thr Leu Asp Asn Gly Lys Ala Ile Ser Ser Ser Arg Gly Asp  
 15 110 115 120

gtt gat tta gtc atc aac tat ttg aaa tct tct gct ggc ttt gct gat 435  
 Val Asp Leu Val Ile Asn Tyr Leu Lys Ser Ser Ala Gly Phe Ala Asp  
 125 130 135

aaa att gat ggt aga atg att gat act ggt aga acc cat ttt tct tac 483  
 20 Lys Ile Asp Gly Arg Met Ile Asp Thr Gly Arg Thr His Phe Ser Tyr  
 140 145 150

act aag aga cag cct ttg ggt gtt tgt ggg cag att att cct tgg aat 531  
 Thr Lys Arg Gln Pro Leu Gly Val Cys Gly Gln Ile Ile Pro Trp Asn  
 155 160 165

25 ttc cca ctg ttg atg tgg gcc tgg aag att gcc cct gct ttg gtc acc 579  
 Phe Pro Leu Leu Met Trp Ala Trp Lys Ile Ala Pro Ala Leu Val Thr  
 170 175 180 185

```

      ggt aac acc gtc gtg ttg aag act gcc gaa tcc acc cca ttg tcc gct   627
      Gly Asn Thr Val Val Leu Lys Thr Ala Glu Ser Thr Pro Leu Ser Ala
                190                195                200

      ttg tat gtg tct aaa tac atc cca cag gcg ggt att cca cct ggt gtg   675
5  Leu Tyr Val Ser Lys Tyr Ile Pro Gln Ala Gly Ile Pro Pro Gly Val
                205                210                215

      atc aac att gta tcc ggg ttt ggt aag att gtg gtt gag gcc att aca   723
      Ile Asn Ile Val Ser Gly Phe Gly Lys Ile Val Val Glu Ala Ile Thr
                220                225                230

10  aac cat cca aaa atc aaa aag gtt gcc ttc aca ggg tcc acg gct acg   771
      Asn His Pro Lys Ile Lys Lys Val Ala Phe Thr Gly Ser Thr Ala Thr
                235                240                245

      ggt aga cac att tac cag tcc gca gcc gca ggc ttg aaa aaa gtg act   819
      Gly Arg His Ile Tyr Gln Ser Ala Ala Ala Gly Leu Lys Lys Val Thr
15  250                255                260                265

      ttg gag ctg ggt ggt aaa tca cca aac att gtc ttc gcg gac gcc gag   867
      Leu Glu Leu Gly Gly Lys Ser Pro Asn Ile Val Phe Ala Asp Ala Glu
                270                275                280

      ttg aaa aaa gcc gtg caa aac att atc ctt ggt atc tac tac aat tct   915
20  Leu Lys Lys Ala Val Gln Asn Ile Ile Leu Gly Ile Tyr Tyr Asn Ser
                285                290                295

      ggt gag gtc tgt tgt gcg ggt tca agg gtg tat gtt gaa gaa tct att   963
      Gly Glu Val Cys Cys Ala Gly Ser Arg Val Tyr Val Glu Glu Ser Ile
                300                305                310

25  tac gac aaa ttc att gaa gag ttc aaa gcc gct tct gaa tcc atc aag   1011
      Tyr Asp Lys Phe Ile Glu Glu Phe Lys Ala Ala Ser Glu Ser Ile Lys
                315                320                325

```



|    |   |      |
|----|---|------|
|    | gtg ggc gac cca ttc gat gaa tct act ttc caa ggt gca caa acc tct | 1059 |
|    | Val Gly Asp Pro Phe Asp Glu Ser Thr Phe Gln Gly Ala Gln Thr Ser |      |
|    | 330 335 340 345   |      |
|    | caa atg caa cta aac aaa atc ttg aaa tac gtt gac att ggt aag aat | 1107 |
| 5  | Gln Met Gln Leu Asn Lys Ile Leu Lys Tyr Val Asp Ile Gly Lys Asn |      |
|    | 350 355 360   |      |
|    | gaa ggt gct act ttg att acc ggt ggt gaa aga tta ggt agc aag ggt | 1155 |
|    | Glu Gly Ala Thr Leu Ile Thr Gly Gly Glu Arg Leu Gly Ser Lys Gly |      |
|    | 365 370 375   |      |
| 10 | tac ttc att aag cca act gtc ttt ggt gac gtt aag gaa gac atg aga | 1203 |
|    | Tyr Phe Ile Lys Pro Thr Val Phe Gly Asp Val Lys Glu Asp Met Arg |      |
|    | 380 385 390   |      |
|    | att gtc aaa gag gaa atc ttt ggc cct gtt gtc act gta acc aaa ttc | 1251 |
|    | Ile Val Lys Glu Glu Ile Phe Gly Pro Val Val Thr Val Thr Lys Phe |      |
| 15 | 395 400 405   |      |
|    | aaa tct gcc gac gaa gtc att aac atg gcg aac gat tct gaa tac ggg | 1299 |
|    | Lys Ser Ala Asp Glu Val Ile Asn Met Ala Asn Asp Ser Glu Tyr Gly |      |
|    | 410 415 420 425   |      |
|    | ttg gct gct ggt att cac acc tct aat att aat acc gcc tta aaa gtg | 1347 |
| 20 | Leu Ala Ala Gly Ile His Thr Ser Asn Ile Asn Thr Ala Leu Lys Val |      |
|    | 430 435 440   |      |
|    | gct gat aga gtt aat gcg ggt acg gtc tgg ata aac act tat aac gat | 1395 |
|    | Ala Asp Arg Val Asn Ala Gly Thr Val Trp Ile Asn Thr Tyr Asn Asp |      |
|    | 445 450 455   |      |
| 25 | ttc cac cac gca gtt cct ttc ggt ggg ttc aat gca tct ggt ttg ggc | 1443 |
|    | Phe His His Ala Val Pro Phe Gly Gly Phe Asn Ala Ser Gly Leu Gly |      |
|    | 460 465 470   |      |

agg gaa atg tct gtt gat gct tta caa aac tac ttg caa gtt aaa gcg 1491  
 Arg Glu Met Ser Val Asp Ala Leu Gln Asn Tyr Leu Gln Val Lys Ala  
 475 480 485

gtc cgt gcc aaa ttg gac gagtaagagc tcgaattcgc 1529  
 5 Val Arg Ala Lys Leu Asp  
 490 495

<210> 2

<211> 495

<212> PRT

10 <213> *Saccharomyces cerevisiae*

<400> 2

Met Ser His Leu Pro Met Thr Val Pro Ile Lys Leu Pro Asn Gly Leu  
 1 5 10 15

Glu Tyr Glu Gln Pro Thr Gly Leu Phe Ile Asn Asn Lys Phe Val Pro  
 15 20 25 30

Ser Lys Gln Asn Lys Thr Phe Glu Val Ile Asn Pro Ser Thr Glu Glu  
 35 40 45

Glu Ile Cys His Ile Tyr Glu Gly Arg Glu Asp Asp Val Glu Glu Ala  
 50 55 60

20 Val Gln Ala Ala Asp Arg Ala Phe Ser Asn Gly Ser Trp Asn Gly Ile  
 65 70 75 80

Asp Pro Ile Asp Arg Gly Lys Ala Leu Tyr Arg Leu Ala Glu Leu Ile  
 85 90 95

Glu Gln Asp Lys Asp Val Ile Ala Ser Ile Glu Thr Leu Asp Asn Gly  
 25 100 105 110

Lys Ala Ile Ser Ser Ser Arg Gly Asp Val Asp Leu Val Ile Asn Tyr  
 115 120 125

Leu Lys Ser Ser Ala Gly Phe Ala Asp Lys Ile Asp Gly Arg Met Ile  
 130 135 140

Asp Thr Gly Arg Thr His Phe Ser Tyr Thr Lys Arg Gln Pro Leu Gly  
 145 150 155 160

5 Val Cys Gly Gln Ile Ile Pro Trp Asn Phe Pro Leu Leu Met Trp Ala  
 165 170 175

Trp Lys Ile Ala Pro Ala Leu Val Thr Gly Asn Thr Val Val Leu Lys  
 180 185 190

Thr Ala Glu Ser Thr Pro Leu Ser Ala Leu Tyr Val Ser Lys Tyr Ile  
 10 195 200 205

Pro Gln Ala Gly Ile Pro Pro Gly Val Ile Asn Ile Val Ser Gly Phe  
 210 215 220

Gly Lys Ile Val Val Glu Ala Ile Thr Asn His Pro Lys Ile Lys Lys  
 225 230 235 240

15 Val Ala Phe Thr Gly Ser Thr Ala Thr Gly Arg His Ile Tyr Gln Ser  
 245 250 255

Ala Ala Ala Gly Leu Lys Lys Val Thr Leu Glu Leu Gly Gly Lys Ser  
 260 265 270

Pro Asn Ile Val Phe Ala Asp Ala Glu Leu Lys Lys Ala Val Gln Asn  
 20 275 280 285

Ile Ile Leu Gly Ile Tyr Tyr Asn Ser Gly Glu Val Cys Cys Ala Gly  
 290 295 300

Ser Arg Val Tyr Val Glu Glu Ser Ile Tyr Asp Lys Phe Ile Glu Glu  
 305 310 315 320

25 Phe Lys Ala Ala Ser Glu Ser Ile Lys Val Gly Asp Pro Phe Asp Glu  
 325 330 335

Ser Thr Phe Gln Gly Ala Gln Thr Ser Gln Met Gln Leu Asn Lys Ile  
340 345 350

Leu Lys Tyr Val Asp Ile Gly Lys Asn Glu Gly Ala Thr Leu Ile Thr  
355 360 365

5 Gly Gly Glu Arg Leu Gly Ser Lys Gly Tyr Phe Ile Lys Pro Thr Val  
370 375 380

Phe Gly Asp Val Lys Glu Asp Met Arg Ile Val Lys Glu Glu Ile Phe  
385 390 395 400

Gly Pro Val Val Thr Val Thr Lys Phe Lys Ser Ala Asp Glu Val Ile  
10 405 410 415

Asn Met Ala Asn Asp Ser Glu Tyr Gly Leu Ala Ala Gly Ile His Thr  
420 425 430

Ser Asn Ile Asn Thr Ala Leu Lys Val Ala Asp Arg Val Asn Ala Gly  
435 440 445

15 Thr Val Trp Ile Asn Thr Tyr Asn Asp Phe His His Ala Val Pro Phe  
450 455 460

Gly Gly Phe Asn Ala Ser Gly Leu Gly Arg Glu Met Ser Val Asp Ala  
465 470 475 480

Leu Gln Asn Tyr Leu Gln Val Lys Ala Val Arg Ala Lys Leu Asp  
20 485 490 495

<210> 3

<211> 1541

<212> DNA

<213> Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (22) .. (1521)

&lt;400&gt; 3

```

5 gcggtaccaa ggagatatca t atg tca gcc gcc gcc acc cag gcc gtg cct 51
      Met Ser Ala Ala Ala Thr Gln Ala Val Pro
              1              5              10

gcc ccc aac cag cag ccc gag gtc ttc tgc aac cag att ttc ata aac 99
Ala Pro Asn Gln Gln Pro Glu Val Phe Cys Asn Gln Ile Phe Ile Asn
10              15              20              25

aat gaa tgg cac gat gcc gtc agc agg aaa aca ttc ccc acc gtc aat 147
Asn Glu Trp His Asp Ala Val Ser Arg Lys Thr Phe Pro Thr Val Asn
              30              35              40

ccg tcc act gga gag gtc atc tgt cag gta gct gaa ggg gac aag gaa 195
15 Pro Ser Thr Gly Glu Val Ile Cys Gln Val Ala Glu Gly Asp Lys Glu
              45              50              55

gat gtg gac aag gca cgt gaa ggc cgc ccg ggc gcc ttc cag ctg ggc 243
Asp Val Asp Lys Ala Arg Glu Gly Arg Pro Gly Ala Phe Gln Leu Gly
              60              65              70

20 tca cct tgg cgc cgc atg gac gca tca cac agc ggc cgg ctg ctg aac 291
Ser Pro Trp Arg Arg Met Asp Ala Ser His Ser Gly Arg Leu Leu Asn
              75              80              85              90

cgc ctg gcc gat ctg atc gag cgg gac cgg acc tac ctg gcg gcc ttg 339
Arg Leu Ala Asp Leu Ile Glu Arg Asp Arg Thr Tyr Leu Ala Ala Leu
25              95              100              105

gag acc ctg gac aat ggc aag ccc tat gtc atc tcc tac ctg gtg gat 387
Glu Thr Leu Asp Asn Gly Lys Pro Tyr Val Ile Ser Tyr Leu Val Asp
              110              115              120

```

|    |   |     |
|----|---|-----|
|    | ttg gac atg gtc ctc aaa tgt ctc cgg tat tat gcc gcc tgg gct gat | 435 |
|    | Leu Asp Met Val Leu Lys Cys Leu Arg Tyr Tyr Ala Gly Trp Ala Asp |     |
|    | 125 130 135   |     |
|    | aag tac cac ggg aaa acc atc ccc att gac gga gac ttc ttc agc tac | 483 |
| 5  | Lys Tyr His Gly Lys Thr Ile Pro Ile Asp Gly Asp Phe Phe Ser Tyr |     |
|    | 140 145 150   |     |
|    | aca cgc cat gaa cct gtg ggg gtg tgc ggg cag atc att ccg tgg aat | 531 |
|    | Thr Arg His Glu Pro Val Gly Val Cys Gly Gln Ile Ile Pro Trp Asn |     |
|    | 155 160 165 170   |     |
| 10 | ttc ccg ctc ctg atg caa gca tgg aag ctg ggc cca gcc ttg gca act | 579 |
|    | Phe Pro Leu Leu Met Gln Ala Trp Lys Leu Gly Pro Ala Leu Ala Thr |     |
|    | 175 180 185   |     |
|    | gga aac gtg gtt gtg atg aag gta gct gag cag aca ccc ctc acc gcc | 627 |
|    | Gly Asn Val Val Val Met Lys Val Ala Glu Gln Thr Pro Leu Thr Ala |     |
| 15 | 190 195 200   |     |
|    | ctc tat gtg gcc aac ctg atc aag gag gct ggc ttt ccc cct ggt gtg | 675 |
|    | Leu Tyr Val Ala Asn Leu Ile Lys Glu Ala Gly Phe Pro Pro Gly Val |     |
|    | 205 210 215   |     |
|    | gtc aac att gtg cct gga ttt ggc ccc acg gct ggg gcc gcc att gcc | 723 |
| 20 | Val Asn Ile Val Pro Gly Phe Gly Pro Thr Ala Gly Ala Ala Ile Ala |     |
|    | 220 225 230   |     |
|    | tcc cat gag gat gtg gac aaa gtg gca ttc aca ggc tcc act gag att | 771 |
|    | Ser His Glu Asp Val Asp Lys Val Ala Phe Thr Gly Ser Thr Glu Ile |     |
|    | 235 240 245 250   |     |
| 25 | ggc cgc gta atc cag gtt gct gct ggg agc agc aac ctc aag aga gtg | 819 |
|    | Gly Arg Val Ile Gln Val Ala Ala Gly Ser Ser Asn Leu Lys Arg Val |     |
|    | 255 260 265   |     |

acc ttg gag ctg ggg ggg aag agc ccc aac atc atc atg tca gat gcc 867  
 Thr Leu Glu Leu Gly Gly Lys Ser Pro Asn Ile Ile Met Ser Asp Ala  
 270 275 280

gat atg gat tgg gcc gtg gaa cag gcc cac ttc gcc ctg ttc ttc aac 915  
 5 Asp Met Asp Trp Ala Val Glu Gln Ala His Phe Ala Leu Phe Phe Asn  
 285 290 295

cag ggc cag tgc tgc tgt gcc ggc tcc cgg acc ttc gtg cag gag gac 963  
 Gln Gly Gln Cys Cys Cys Ala Gly Ser Arg Thr Phe Val Gln Glu Asp  
 300 305 310

10 atc tat gat gag ttt gtg gtg cgg agc gtt gcc cgg gcc aag tct cgg 1011  
 Ile Tyr Asp Glu Phe Val Val Arg Ser Val Ala Arg Ala Lys Ser Arg  
 315 320 325 330

gtg gtc ggg aac ccc ttt gat agc aag acc gag cag ggg ccg cag gtg 1059  
 Val Val Gly Asn Pro Phe Asp Ser Lys Thr Glu Gln Gly Pro Gln Val  
 15 335 340 345

gat gaa act cag ttt aag aag atc ctc ggc tac atc aac acg ggg aag 1107  
 Asp Glu Thr Gln Phe Lys Lys Ile Leu Gly Tyr Ile Asn Thr Gly Lys  
 350 355 360

caa gag ggg gcg aag ctg ctg tgt ggt ggg ggc att gct gct gac cgt 1155  
 20 Gln Glu Gly Ala Lys Leu Leu Cys Gly Gly Gly Ile Ala Ala Asp Arg  
 365 370 375

ggt tac ttc atc cag ccc act gtg ttt gga gat gtg cag gat ggc atg 1203  
 Gly Tyr Phe Ile Gln Pro Thr Val Phe Gly Asp Val Gln Asp Gly Met  
 380 385 390

25 acc atc gcc aag gag gag atc ttc ggg cca gtg atg cag atc ctg aag 1251  
 Thr Ile Ala Lys Glu Glu Ile Phe Gly Pro Val Met Gln Ile Leu Lys  
 395 400 405 410

ttc aag acc ata gag gag gtt gtt ggg aga gcc aac aat tcc acg tac 1299  
 Phe Lys Thr Ile Glu Glu Val Val Gly Arg Ala Asn Asn Ser Thr Tyr  
 415 420 425

ggg ctg gcc gca gct gtc ttc aca aag gat ttg gac aag gcc aat tac 1347  
 5 Gly Leu Ala Ala Ala Val Phe Thr Lys Asp Leu Asp Lys Ala Asn Tyr  
 430 435 440

ctg tcc cag gcc ctc cag gcg ggc act gtg tgg gtc aac tgc tat gat 1395  
 Leu Ser Gln Ala Leu Gln Ala Gly Thr Val Trp Val Asn Cys Tyr Asp  
 445 450 455

10 gtg ttt gga gcc cag tca ccc ttt ggt ggc tac aag atg tcg ggg agt 1443  
 Val Phe Gly Ala Gln Ser Pro Phe Gly Gly Tyr Lys Met Ser Gly Ser  
 460 465 470

ggc cgg gag ttg ggc gag tac ggg ctg cag gca tac act gaa gtg aaa 1491  
 Gly Arg Glu Leu Gly Glu Tyr Gly Leu Gln Ala Tyr Thr Glu Val Lys  
 15 475 480 485 490

act gtc aca gtc aaa gtg cct cag aag aac tcataagagc tcgaattcgc 1541  
 Thr Val Thr Val Lys Val Pro Gln Lys Asn  
 495 500

<210> 4  
 20 <211> 500  
 <212> PRT  
 <213> Homo sapiens

<400> 4  
 Met Ser Ala Ala Ala Thr Gln Ala Val Pro Ala Pro Asn Gln Gln Pro  
 25 1 5 10 15  
 Glu Val Phe Cys Asn Gln Ile Phe Ile Asn Asn Glu Trp His Asp Ala  
 20 25 30



Val Ser Arg Lys Thr Phe Pro Thr Val Asn Pro Ser Thr Gly Glu Val  
35 40 45

Ile Cys Gln Val Ala Glu Gly Asp Lys Glu Asp Val Asp Lys Ala Arg  
50 55 60

5 Glu Gly Arg Pro Gly Ala Phe Gln Leu Gly Ser Pro Trp Arg Arg Met  
65 70 75 80

Asp Ala Ser His Ser Gly Arg Leu Leu Asn Arg Leu Ala Asp Leu Ile  
85 90 95

Glu Arg Asp Arg Thr Tyr Leu Ala Ala Leu Glu Thr Leu Asp Asn Gly  
10 100 105 110

Lys Pro Tyr Val Ile Ser Tyr Leu Val Asp Leu Asp Met Val Leu Lys  
115 120 125

Cys Leu Arg Tyr Tyr Ala Gly Trp Ala Asp Lys Tyr His Gly Lys Thr  
130 135 140

15 Ile Pro Ile Asp Gly Asp Phe Phe Ser Tyr Thr Arg His Glu Pro Val  
145 150 155 160

Gly Val Cys Gly Gln Ile Ile Pro Trp Asn Phe Pro Leu Leu Met Gln  
165 170 175

Ala Trp Lys Leu Gly Pro Ala Leu Ala Thr Gly Asn Val Val Val Met  
20 180 185 190

Lys Val Ala Glu Gln Thr Pro Leu Thr Ala Leu Tyr Val Ala Asn Leu  
195 200 205

Ile Lys Glu Ala Gly Phe Pro Pro Gly Val Val Asn Ile Val Pro Gly  
210 215 220

Phe Gly Pro Thr Ala Gly Ala Ala Ile Ala Ser His Glu Asp Val Asp  
225 230 235 240

Lys Val Ala Phe Thr Gly Ser Thr Glu Ile Gly Arg Val Ile Gln Val  
245 250 255

5 Ala Ala Gly Ser Ser Asn Leu Lys Arg Val Thr Leu Glu Leu Gly Gly  
260 265 270

Lys Ser Pro Asn Ile Ile Met Ser Asp Ala Asp Met Asp Trp Ala Val  
275 280 285

Glu Gln Ala His Phe Ala Leu Phe Phe Asn Gln Gly Gln Cys Cys Cys  
10 290 295 300

Ala Gly Ser Arg Thr Phe Val Gln Glu Asp Ile Tyr Asp Glu Phe Val  
305 310 315 320

Val Arg Ser Val Ala Arg Ala Lys Ser Arg Val Val Gly Asn Pro Phe  
325 330 335

15 Asp Ser Lys Thr Glu Gln Gly Pro Gln Val Asp Glu Thr Gln Phe Lys  
340 345 350

Lys Ile Leu Gly Tyr Ile Asn Thr Gly Lys Gln Glu Gly Ala Lys Leu  
355 360 365

Leu Cys Gly Gly Gly Ile Ala Ala Asp Arg Gly Tyr Phe Ile Gln Pro  
20 370 375 380

Thr Val Phe Gly Asp Val Gln Asp Gly Met Thr Ile Ala Lys Glu Glu  
385 390 395 400

Ile Phe Gly Pro Val Met Gln Ile Leu Lys Phe Lys Thr Ile Glu Glu  
405 410 415

Val Val Gly Arg Ala Asn Asn Ser Thr Tyr Gly Leu Ala Ala Ala Val  
 420 425 430

Phe Thr Lys Asp Leu Asp Lys Ala Asn Tyr Leu Ser Gln Ala Leu Gln  
 435 440 445

5 Ala Gly Thr Val Trp Val Asn Cys Tyr Asp Val Phe Gly Ala Gln Ser  
 450 455 460

Pro Phe Gly Gly Tyr Lys Met Ser Gly Ser Gly Arg Glu Leu Gly Glu  
 465 470 475 480

Tyr Gly Leu Gln Ala Tyr Thr Glu Val Lys Thr Val Thr Val Lys Val  
 10 485 490 495

Pro Gln Lys Asn  
 500

<210> 5  
 <211> 1512  
 15 <212> DNA  
 <213> Escherichia coli

<220>  
 <221> CDS  
 <222> (37) .. (1473)

20 <400> 5  
 gctaccatgg cttaaccggt accaaggaga tatcat atg tca gta ccc gtt caa 54  
 Met Ser Val Pro Val Gln  
 1 5

cat cct atg tat atc gat gga cag ttt gtt acc tgg cgt gga gac gca 102  
 25 His Pro Met Tyr Ile Asp Gly Gln Phe Val Thr Trp Arg Gly Asp Ala  
 10 15 20

|    |   |     |
|----|---|-----|
|    | tgg att gat gtg gta aac cct gct aca gag gct gtc att tcc cgc ata | 150 |
|    | Trp Ile Asp Val Val Asn Pro Ala Thr Glu Ala Val Ile Ser Arg Ile |     |
|    | 25 30 35  |     |
|    | ccc gat ggt cag gcc gag gat gcc cgt aag gca atc gat gca gca gaa | 198 |
| 5  | Pro Asp Gly Gln Ala Glu Asp Ala Arg Lys Ala Ile Asp Ala Ala Glu |     |
|    | 40 45 50  |     |
|    | cgt gca caa cca gaa tgg gaa gcg ttg cct gct att gaa cgc gcc agt | 246 |
|    | Arg Ala Gln Pro Glu Trp Glu Ala Leu Pro Ala Ile Glu Arg Ala Ser |     |
|    | 55 60 65 70   |     |
| 10 | tgg ttg cgc aaa atc tcc gcc ggg atc cgc gaa cgc gcc agt gaa atc | 294 |
|    | Trp Leu Arg Lys Ile Ser Ala Gly Ile Arg Glu Arg Ala Ser Glu Ile |     |
|    | 75 80 85  |     |
|    | agt gcg ctg att gtt gaa gaa ggg ggc aag atc cag cag ctg gct gaa | 342 |
|    | Ser Ala Leu Ile Val Glu Glu Gly Gly Lys Ile Gln Gln Leu Ala Glu |     |
| 15 | 90 95 100   |     |
|    | gtc gaa gtg gct ttt act gcc gac tat atc gat tac atg gcg gag tgg | 390 |
|    | Val Glu Val Ala Phe Thr Ala Asp Tyr Ile Asp Tyr Met Ala Glu Trp |     |
|    | 105 110 115   |     |
|    | gca cgg cgt tac gag ggc gag att att caa agc gat cgt cca gga gaa | 438 |
| 20 | Ala Arg Arg Tyr Glu Gly Glu Ile Ile Gln Ser Asp Arg Pro Gly Glu |     |
|    | 120 125 130   |     |
|    | aat att ctt ttg ttt aaa cgt gcg ctt ggt gtg act acc ggc att ctg | 486 |
|    | Asn Ile Leu Leu Phe Lys Arg Ala Leu Gly Val Thr Thr Gly Ile Leu |     |
|    | 135 140 145 150   |     |
| 25 | ccg tgg aac ttc ccg ttc ttc ctc att gcc cgc aaa atg gct ccc gct | 534 |
|    | Pro Trp Asn Phe Pro Phe Phe Leu Ile Ala Arg Lys Met Ala Pro Ala |     |
|    | 155 160 165   |     |

ctt ttg acc ggt aat acc atc gtc att aaa cct agt gaa ttt acg aca 582  
 Leu Leu Thr Gly Asn Thr Ile Val Ile Lys Pro Ser Glu Phe Thr Thr  
 170 175 180

aac aat gcg att gca ttc gcc aaa atc gtc gat gaa ata ggc ctt ccg 630  
 5 Asn Asn Ala Ile Ala Phe Ala Lys Ile Val Asp Glu Ile Gly Leu Pro  
 185 190 195

cgc ggc gtg ttt aac ctt gta ctg ggg cgt ggt gaa acc gtt ggg caa 678  
 Arg Gly Val Phe Asn Leu Val Leu Gly Arg Gly Glu Thr Val Gly Gln  
 200 205 210

10 gaa ctg gcg ggt aac cca aag gtc gca atg gtc agt atg aca ggc agc 726  
 Glu Leu Ala Gly Asn Pro Lys Val Ala Met Val Ser Met Thr Gly Ser  
 215 220 225 230

gtc tct gca ggt gag aag atc atg gcg act gcg gcg aaa aac atc acc 774  
 Val Ser Ala Gly Glu Lys Ile Met Ala Thr Ala Ala Lys Asn Ile Thr  
 15 235 240 245

aaa gtg tgt ctg gaa ttg ggg ggt aaa gca cca gct atc gta atg gac 822  
 Lys Val Cys Leu Glu Leu Gly Gly Lys Ala Pro Ala Ile Val Met Asp  
 250 255 260

gat gcc gat ctt gaa ctg gca gtc aaa gcc atc gtt gat tca cgc gtc 870  
 20 Asp Ala Asp Leu Glu Leu Ala Val Lys Ala Ile Val Asp Ser Arg Val  
 265 270 275

att aat agt ggg caa gtg tgt aac tgt gca gaa cgt gtt tat gta cag 918  
 Ile Asn Ser Gly Gln Val Cys Asn Cys Ala Glu Arg Val Tyr Val Gln  
 280 285 290

25 aaa ggc att tat gat cag ttc gtc aat cgg ctg ggt gaa gcg atg cag 966  
 Lys Gly Ile Tyr Asp Gln Phe Val Asn Arg Leu Gly Glu Ala Met Gln  
 295 300 305 310

gcg gtt caa ttt ggt aac ccc gct gaa cgc aac gac att gcg atg ggg 1014  
 Ala Val Gln Phe Gly Asn Pro Ala Glu Arg Asn Asp Ile Ala Met Gly  
 315 320 325

ccg ttg att aac gcc gcg gcg ctg gaa agg gtc gag caa aaa gtg gcg 1062  
 5 Pro Leu Ile Asn Ala Ala Ala Leu Glu Arg Val Glu Gln Lys Val Ala  
 330 335 340

cgc gca gta gaa gaa ggg gcg aga gtg gcg ttc ggt ggc aaa gcg gta 1110  
 Arg Ala Val Glu Glu Gly Ala Arg Val Ala Phe Gly Gly Lys Ala Val  
 345 350 355

10 gag ggg aaa gga tat tat tat ccg ccg aca ttg ctg ctg gat gtt cgc 1158  
 Glu Gly Lys Gly Tyr Tyr Tyr Pro Pro Thr Leu Leu Leu Asp Val Arg  
 360 365 370

cag gaa atg tcg att atg cat gag gaa acc ttt ggc ccg gtg ctg cca 1206  
 Gln Glu Met Ser Ile Met His Glu Glu Thr Phe Gly Pro Val Leu Pro  
 15 375 380 385 390

gtt gtc gca ttt gac acg ctg gaa gat gct atc tca atg gct aat gac 1254  
 Val Val Ala Phe Asp Thr Leu Glu Asp Ala Ile Ser Met Ala Asn Asp  
 395 400 405

agt gat tac ggc ctg acc tca tca atc tat acc caa aat ctg aac gtc 1302  
 20 Ser Asp Tyr Gly Leu Thr Ser Ser Ile Tyr Thr Gln Asn Leu Asn Val  
 410 415 420

gcg atg aaa gcc att aaa ggg ctg aag ttt ggt gaa act tac atc aac 1350  
 Ala Met Lys Ala Ile Lys Gly Leu Lys Phe Gly Glu Thr Tyr Ile Asn  
 425 430 435

25 cgt gaa aac ttc gaa gct atg caa ggc ttc cac gcc gga tgg cgt aaa 1398  
 Arg Glu Asn Phe Glu Ala Met Gln Gly Phe His Ala Gly Trp Arg Lys  
 440 445 450

tcc ggt att ggc ggc gca gat ggt aaa cat ggc ttg cat gga tat ctg 1446  
 Ser Gly Ile Gly Gly Ala Asp Gly Lys His Gly Leu His Gly Tyr Leu  
 455 460 465 470

cag acc cag gtg gtt tat tta cag tct taagagctcg aattcccgtc 1493  
 5 Gln Thr Gln Val Val Tyr Leu Gln Ser  
 475

gacggctcta gactcgagcg 1513

<210> 6  
 <211> 479  
 10 <212> PRT  
 <213> Escherichia coli

<400> 6  
 Met Ser Val Pro Val Gln His Pro Met Tyr Ile Asp Gly Gln Phe Val  
 1 5 10 15

15 Thr Trp Arg Gly Asp Ala Trp Ile Asp Val Val Asn Pro Ala Thr Glu  
 20 25 30

Ala Val Ile Ser Arg Ile Pro Asp Gly Gln Ala Glu Asp Ala Arg Lys  
 35 40 45

Ala Ile Asp Ala Ala Glu Arg Ala Gln Pro Glu Trp Glu Ala Leu Pro  
 20 50 55 60

Ala Ile Glu Arg Ala Ser Trp Leu Arg Lys Ile Ser Ala Gly Ile Arg  
 65 70 75 80

Glu Arg Ala Ser Glu Ile Ser Ala Leu Ile Val Glu Glu Gly Gly Lys  
 85 90 95

25 Ile Gln Gln Leu Ala Glu Val Glu Val Ala Phe Thr Ala Asp Tyr Ile  
 100 105 110

Asp Tyr Met Ala Glu Trp Ala Arg Arg Tyr Glu Gly Glu Ile Ile Gln  
115 120 125

Ser Asp Arg Pro Gly Glu Asn Ile Leu Leu Phe Lys Arg Ala Leu Gly  
130 135 140

5 Val Thr Thr Gly Ile Leu Pro Trp Asn Phe Pro Phe Phe Leu Ile Ala  
145 150 155 160

Arg Lys Met Ala Pro Ala Leu Leu Thr Gly Asn Thr Ile Val Ile Lys  
165 170 175

Pro Ser Glu Phe Thr Thr Asn Asn Ala Ile Ala Phe Ala Lys Ile Val  
10 180 185 190

Asp Glu Ile Gly Leu Pro Arg Gly Val Phe Asn Leu Val Leu Gly Arg  
195 200 205

Gly Glu Thr Val Gly Gln Glu Leu Ala Gly Asn Pro Lys Val Ala Met  
210 215 220

15 Val Ser Met Thr Gly Ser Val Ser Ala Gly Glu Lys Ile Met Ala Thr  
225 230 235 240

Ala Ala Lys Asn Ile Thr Lys Val Cys Leu Glu Leu Gly Gly Lys Ala  
245 250 255

Pro Ala Ile Val Met Asp Asp Ala Asp Leu Glu Leu Ala Val Lys Ala  
20 260 265 270

Ile Val Asp Ser Arg Val Ile Asn Ser Gly Gln Val Cys Asn Cys Ala  
275 280 285

Glu Arg Val Tyr Val Gln Lys Gly Ile Tyr Asp Gln Phe Val Asn Arg  
290 295 300



Leu Gly Glu Ala Met Gln Ala Val Gln Phe Gly Asn Pro Ala Glu Arg  
305 310 315 320

Asn Asp Ile Ala Met Gly Pro Leu Ile Asn Ala Ala Ala Leu Glu Arg  
325 330 335

5 Val Glu Gln Lys Val Ala Arg Ala Val Glu Glu Gly Ala Arg Val Ala  
340 345 350

Phe Gly Gly Lys Ala Val Glu Gly Lys Gly Tyr Tyr Tyr Pro Pro Thr  
355 360 365

Leu Leu Leu Asp Val Arg Gln Glu Met Ser Ile Met His Glu Glu Thr  
10 370 375 380

Phe Gly Pro Val Leu Pro Val Val Ala Phe Asp Thr Leu Glu Asp Ala  
385 390 395 400

Ile Ser Met Ala Asn Asp Ser Asp Tyr Gly Leu Thr Ser Ser Ile Tyr  
405 410 415

15 Thr Gln Asn Leu Asn Val Ala Met Lys Ala Ile Lys Gly Leu Lys Phe  
420 425 430

Gly Glu Thr Tyr Ile Asn Arg Glu Asn Phe Glu Ala Met Gln Gly Phe  
435 440 445

His Ala Gly Trp Arg Lys Ser Gly Ile Gly Gly Ala Asp Gly Lys His  
20 450 455 460

Gly Leu His Gly Tyr Leu Gln Thr Gln Val Val Tyr Leu Gln Ser  
465 470 475

<210> 7  
 <211> 1574  
 <212> DNA  
 <213> Escherichia coli

5 <220>  
 <221> CDS  
 <222> (22)..(1557)

<400> 7

|    |   |     |
|----|---|-----|
| 10 | gcggtaccaa ggaggtatca t atg acc aat aat ccc cct tca gca cag att | 51  |
|    | Met Thr Asn Asn Pro Pro Ser Ala Gln Ile                         |     |
|    | 1 5 10  |     |
|    | aag ccc ggc gag tat ggt ttc ccc ctc aag tta aaa gcc cgc tat gac | 99  |
|    | Lys Pro Gly Glu Tyr Gly Phe Pro Leu Lys Leu Lys Ala Arg Tyr Asp |     |
|    | 15 20 25  |     |
| 15 | aac ttt att ggc ggc gaa tgg gta gcc cct gcc gac ggc gag tat tac | 147 |
|    | Asn Phe Ile Gly Gly Glu Trp Val Ala Pro Ala Asp Gly Glu Tyr Tyr |     |
|    | 30 35 40  |     |
| 20 | cag aat ctg acg ccg gtg acc ggg cag ctg ctg tgc gaa gtg gcg tct | 195 |
|    | Gln Asn Leu Thr Pro Val Thr Gly Gln Leu Leu Cys Glu Val Ala Ser |     |
|    | 45 50 55  |     |
|    | tcg ggc aaa cga gac atc gat ctg gcg ctg gat gct gcg cac aaa gtg | 243 |
|    | Ser Gly Lys Arg Asp Ile Asp Leu Ala Leu Asp Ala Ala His Lys Val |     |
|    | 60 65 70  |     |
| 25 | aaa gat aaa tgg gcg cac acc tcg gtg cag gat cgt gcg gcg att ctg | 291 |
|    | Lys Asp Lys Trp Ala His Thr Ser Val Gln Asp Arg Ala Ala Ile Leu |     |
|    | 75 80 85 90   |     |
|    | ttt aag att gcc gat cga atg gaa caa aac ctc gag ctg tta gcg aca | 339 |
|    | Phe Lys Ile Ala Asp Arg Met Glu Gln Asn Leu Glu Leu Leu Ala Thr |     |
|    | 95 100 105  |     |

|    |   |     |
|----|---|-----|
|    | gct gaa acc tgg gat aac ggc aaa ccc att cgc gaa acc agt gct gcg | 387 |
|    | Ala Glu Thr Trp Asp Asn Gly Lys Pro Ile Arg Glu Thr Ser Ala Ala |     |
|    | 110 115 120   |     |
|    | gat gta ccg ctg gcg att gac cat ttc cgc tat ttc gcc tcg tgt att | 435 |
| 5  | Asp Val Pro Leu Ala Ile Asp His Phe Arg Tyr Phe Ala Ser Cys Ile |     |
|    | 125 130 135   |     |
|    | cgg gcg cag gaa ggt ggg atc agt gaa gtt gat agc gaa acc gtg gcc | 483 |
|    | Arg Ala Gln Glu Gly Gly Ile Ser Glu Val Asp Ser Glu Thr Val Ala |     |
|    | 140 145 150   |     |
| 10 | tat cat ttc cat gaa ccg tta ggc gtg gtg ggg cag att atc ccg tgg | 531 |
|    | Tyr His Phe His Glu Pro Leu Gly Val Val Gly Gln Ile Ile Pro Trp |     |
|    | 155 160 165 170   |     |
|    | aac ttc ccg ctg ctg atg gcg agc tgg aaa atg gct ccc gcg ctg gcg | 579 |
|    | Asn Phe Pro Leu Leu Met Ala Ser Trp Lys Met Ala Pro Ala Leu Ala |     |
| 15 | 175 180 185   |     |
|    | gcg ggc aac tgt gtg gtg ctg aaa ccc gca cgt ctt acc ccg ctt tct | 627 |
|    | Ala Gly Asn Cys Val Val Leu Lys Pro Ala Arg Leu Thr Pro Leu Ser |     |
|    | 190 195 200   |     |
|    | gta ctg ctg cta atg gaa att gtc ggt gat tta ctg ccg ccg ggc gtg | 675 |
| 20 | Val Leu Leu Leu Met Glu Ile Val Gly Asp Leu Leu Pro Pro Gly Val |     |
|    | 205 210 215   |     |
|    | gtg aac gtg gtc aat ggc gca ggt ggg gta att ggc gaa tat ctg gcg | 723 |
|    | Val Asn Val Val Asn Gly Ala Gly Gly Val Ile Gly Glu Tyr Leu Ala |     |
|    | 220 225 230   |     |
| 25 | acc tcg aaa cgc atc gcc aaa gtg gcg ttt acc ggc tca acg gaa gtg | 771 |
|    | Thr Ser Lys Arg Ile Ala Lys Val Ala Phe Thr Gly Ser Thr Glu Val |     |
|    | 235 240 245 250   |     |

ggc caa caa att atg caa tac gca acg caa aac att att ccg gtg acg 819  
 Gly Gln Gln Ile Met Gln Tyr Ala Thr Gln Asn Ile Ile Pro Val Thr  
 255 260 265

ctg gag ttg ggc ggt aag tcg cca aat atc gtc ttt gct gat gtg atg 867  
 5 Leu Glu Leu Gly Gly Lys Ser Pro Asn Ile Val Phe Ala Asp Val Met  
 270 275 280

gat gaa gaa gat gcc ttt ttc gat aaa gcg ctg gaa ggc ttt gca ctg 915  
 Asp Glu Glu Asp Ala Phe Phe Asp Lys Ala Leu Glu Gly Phe Ala Leu  
 285 290 295

10 ttt gcc ttt aac cag ggc gaa gtt tgc acc tgt ccg agt cgt gct tta 963  
 Phe Ala Phe Asn Gln Gly Glu Val Cys Thr Cys Pro Ser Arg Ala Leu  
 300 305 310

gtg cag gaa tct atc tac gaa cgc ttt atg gaa cgc gcc atc cgc cgt 1011  
 Val Gln Glu Ser Ile Tyr Glu Arg Phe Met Glu Arg Ala Ile Arg Arg  
 15 315 320 325 330

gtc gaa agc att cgt agc ggt aac ccg ctc gac agc gtg acg caa atg 1059  
 Val Glu Ser Ile Arg Ser Gly Asn Pro Leu Asp Ser Val Thr Gln Met  
 335 340 345

ggc gcg cag gtt tct cac ggg caa ctg gaa acc atc ctc aac tac att 1107  
 20 Gly Ala Gln Val Ser His Gly Gln Leu Glu Thr Ile Leu Asn Tyr Ile  
 350 355 360

gat atc ggt aaa aaa gag ggc gct gac gtg ctc aca ggc ggg cgg cgc 1155  
 Asp Ile Gly Lys Lys Glu Gly Ala Asp Val Leu Thr Gly Gly Arg Arg  
 365 370 375

25 aag ctg ctg gaa ggt gaa ctg aaa gac ggc tac tac ctc gaa ccg acg 1203  
 Lys Leu Leu Glu Gly Glu Leu Lys Asp Gly Tyr Tyr Leu Glu Pro Thr  
 380 385 390

|    |  |      |
|----|--|------|
|    | att ctg ttt ggt cag aac aat atg cgg gtg ttc cag gag gag att ttt                | 1251 |
|    | Ile Leu Phe Gly Gln Asn Asn Met Arg Val Phe Gln Glu Glu Ile Phe                |      |
|    | 395                      400                      405                      410 |      |
|    | ggc ccg gtg ctg gcg gtg acc acc ttc aaa acg atg gaa gaa gcg ctg                | 1299 |
| 5  | Gly Pro Val Leu Ala Val Thr Thr Phe Lys Thr Met Glu Glu Ala Leu                |      |
|    | 415                      420                      425                          |      |
|    | gag ctg gcg aac gat acg caa tat ggc ctg ggc gcg ggc gtc tgg agc                | 1347 |
|    | Glu Leu Ala Asn Asp Thr Gln Tyr Gly Leu Gly Ala Gly Val Trp Ser                |      |
|    | 430                      435                      440                          |      |
| 10 | cgc aac ggt aat ctg gcc tat aag atg ggg cgc ggc ata cag gct ggg                | 1395 |
|    | Arg Asn Gly Asn Leu Ala Tyr Lys Met Gly Arg Gly Ile Gln Ala Gly                |      |
|    | 445                      450                      455                          |      |
|    | cgc gtg tgg acc aac tgt tat cac gct tac ccg gca cat gcg gcg ttt                | 1443 |
|    | Arg Val Trp Thr Asn Cys Tyr His Ala Tyr Pro Ala His Ala Ala Phe                |      |
| 15 | 460                      465                      470                          |      |
|    | ggt ggc tac aaa caa tca ggt atc ggt cgc gaa acc cac aag atg atg                | 1491 |
|    | Gly Gly Tyr Lys Gln Ser Gly Ile Gly Arg Glu Thr His Lys Met Met                |      |
|    | 475                      480                      485                      490 |      |
|    | ctg gag cat tac cag caa acc aag tgc ctg ctg gtg agc tac tcg gat                | 1539 |
| 20 | Leu Glu His Tyr Gln Gln Thr Lys Cys Leu Leu Val Ser Tyr Ser Asp                |      |
|    | 495                      500                      505                          |      |
|    | aaa ccg ttg ggg ctg ttc taagagctcg aattcgc                                     | 1574 |
|    | Lys Pro Leu Gly Leu Phe  |      |
|    | 510  |      |

&lt;210&gt; 8

&lt;211&gt; 512

&lt;212&gt; PRT

&lt;213&gt; Escherichia coli

5 &lt;400&gt; 8

Met Thr Asn Asn Pro Pro Ser Ala Gln Ile Lys Pro Gly Glu Tyr Gly

1 5 10 15

Phe Pro Leu Lys Leu Lys Ala Arg Tyr Asp Asn Phe Ile Gly Gly Glu

20 25 30

10 Trp Val Ala Pro Ala Asp Gly Glu Tyr Tyr Gln Asn Leu Thr Pro Val

35 40 45

Thr Gly Gln Leu Leu Cys Glu Val Ala Ser Ser Gly Lys Arg Asp Ile

50 55 60

Asp Leu Ala Leu Asp Ala Ala His Lys Val Lys Asp Lys Trp Ala His

15 65 70 75 80

Thr Ser Val Gln Asp Arg Ala Ala Ile Leu Phe Lys Ile Ala Asp Arg

85 90 95

Met Glu Gln Asn Leu Glu Leu Leu Ala Thr Ala Glu Thr Trp Asp Asn

100 105 110

20 Gly Lys Pro Ile Arg Glu Thr Ser Ala Ala Asp Val Pro Leu Ala Ile

115 120 125

Asp His Phe Arg Tyr Phe Ala Ser Cys Ile Arg Ala Gln Glu Gly Gly

130 135 140

Ile Ser Glu Val Asp Ser Glu Thr Val Ala Tyr His Phe His Glu Pro

25 145 150 155 160

Leu Gly Val Val Gly Gln Ile Ile Pro Trp Asn Phe Pro Leu Leu Met  
 165 170 175

Ala Ser Trp Lys Met Ala Pro Ala Leu Ala Ala Gly Asn Cys Val Val  
 180 185 190

5 Leu Lys Pro Ala Arg Leu Thr Pro Leu Ser Val Leu Leu Leu Met Glu  
 195 200 205

Ile Val Gly Asp Leu Leu Pro Pro Gly Val Val Asn Val Val Asn Gly  
 210 215 220

Ala Gly Gly Val Ile Gly Glu Tyr Leu Ala Thr Ser Lys Arg Ile Ala  
 10 225 230 235 240

Lys Val Ala Phe Thr Gly Ser Thr Glu Val Gly Gln Gln Ile Met Gln  
 245 250 255

Tyr Ala Thr Gln Asn Ile Ile Pro Val Thr Leu Glu Leu Gly Gly Lys  
 260 265 270

15 Ser Pro Asn Ile Val Phe Ala Asp Val Met Asp Glu Glu Asp Ala Phe  
 275 280 285

Phe Asp Lys Ala Leu Glu Gly Phe Ala Leu Phe Ala Phe Asn Gln Gly  
 290 295 300

Glu Val Cys Thr Cys Pro Ser Arg Ala Leu Val Gln Glu Ser Ile Tyr  
 20 305 310 315 320

Glu Arg Phe Met Glu Arg Ala Ile Arg Arg Val Glu Ser Ile Arg Ser  
 325 330 335

Gly Asn Pro Leu Asp Ser Val Thr Gln Met Gly Ala Gln Val Ser His  
 340 345 350

Gly Gln Leu Glu Thr Ile Leu Asn Tyr Ile Asp Ile Gly Lys Lys Glu  
 355 360 365

Gly Ala Asp Val Leu Thr Gly Gly Arg Arg Lys Leu Leu Glu Gly Glu  
 370 375 380

5 Leu Lys Asp Gly Tyr Tyr Leu Glu Pro Thr Ile Leu Phe Gly Gln Asn  
 385 390 395 400

Asn Met Arg Val Phe Gln Glu Glu Ile Phe Gly Pro Val Leu Ala Val  
 405 410 415

10 Thr Thr Phe Lys Thr Met Glu Glu Ala Leu Glu Leu Ala Asn Asp Thr  
 420 425 430

Gln Tyr Gly Leu Gly Ala Gly Val Trp Ser Arg Asn Gly Asn Leu Ala  
 435 440 445

Tyr Lys Met Gly Arg Gly Ile Gln Ala Gly Arg Val Trp Thr Asn Cys  
 450 455 460

15 Tyr His Ala Tyr Pro Ala His Ala Ala Phe Gly Gly Tyr Lys Gln Ser  
 465 470 475 480

Gly Ile Gly Arg Glu Thr His Lys Met Met Leu Glu His Tyr Gln Gln  
 485 490 495

20 Thr Lys Cys Leu Leu Val Ser Tyr Ser Asp Lys Pro Leu Gly Leu Phe  
 500 505 510

&lt;210&gt; 9

&lt;211&gt; 5267

&lt;212&gt; DNA

&lt;213&gt; Klebsiella pneumoniae



&lt;220&gt;

&lt;223&gt; Location complement 300..2153

&lt;220&gt;

&lt;223&gt; Location complement 2166..2591

5 &lt;220&gt;

&lt;223&gt; Location complement 2594..3034

&lt;220&gt;

&lt;223&gt; Location complement 2191..4858

&lt;400&gt; 9

10 agcgctatat gcggtgatgc aatttctatg cgcacccgtt ctcggagcac tgtccgaccg 60  
 ctttggccgc cgcccagtc tgcctgcttc gctacttggg gccactatcg actacgcgat 120  
 catggcgacc acaccggtcc tgtggatctc ccactgacca aagctggccc cgggcgaccg 180  
 cagcgagcgc acgcagcccg cgccgaagaa aatgagcaat ccggtgcca gaaactcggc 240  
 cacgcactgc ccggttaagg tagaagtctg gttcattatc ggcatcctga aatagcacgt 300  
 15 taaagagaga ggctggcgcg agcgcccggt taattcgctt gaccggccag tagcagcccg 360  
 gtggcgaccg cattgcgcgg cccttctgtt ccccgaaat tgccctgccc ggcgaccacg 420  
 ccatagtgcg acaaggcttc cgtgataagc tgcgggatct caaagtccag cgatgagccg 480  
 cccaccagca ccacaaaggc gatatcgcg atggaaccgc cgggtgagac ctggcgagc 540  
 ggcgcgagc agttggtgac aaacactttc tctttcgctt gccggcgcac gagacgaatt 600  
 20 ttttccagcg ggctggcggt atcgatcggc accagttcgc cctccttgat gtacaccact 660  
 ttggcgaaac ccgcccgggt gagggcttcc cgaaagaact ccaccgcgcc attctcgtga 720  
 cgaatactga acaggctttc cactttggcc agcgggtatt tttttatcgc ttccgccagc 780  
 gaaagatcct cgaggcccag ctcggtttta atcaacaggc tgaccatatt ccccgccccg 840  
 gcgagatgga ccgcccgtat ctgcccctcc gcgttgacga tcgccgcac cgtcgagccg 900  
 25 gcgccgaggt cgaggatcgc cagcgccgcc gcacagccgg gagtgggtaa cgccccggcg 960  
 atggccatgt tggcctccac gccgcccacc accacctcgg tctgcagtcg ggcgctcagt 1020  
 tcgggggcga taacctgcat ttgcagacga tccgctttca ccacgcccgc catcccagc 1080  
 gcattctcca tggcgactc gccggccatc ccgcccgtca ccttgcgcg aataaacgta 1140  
 tccaccgcca gcagatcctg gatgtatata gcgctcatct catggccggg cagggaacgc 1200  
 30 attaccttgc gcaccgctc aagcatgcc cgggcggtgg tgcgggttc gccgcggatg 1260  
 tcggttaccg gagcgagcc gctcatcgcc tgcagatgg cttccgcgcc ctcggcgaca 1320  
 tcggcctctc cgccggcgtt ttcgcccgt atgtagaggt tgcggccgg gatcacccgc 1380  
 gactgcacat cccctgccc ggctctgagc accaccgccc aacgggtgcc aatcagggcg 1440

cgggcgatgg ggacgatggc ctgggtctct tccgggctta gcccgaagaa ggtggcgatc 1500  
ccgtagggat tgcacaggat ccgcaccacc tggcccggcg cgccacttc caccgccgcc 1560  
attacccctt cggggacctg ctccagcagc gtcacttcat ccaccaccgg cagggtttta 1620  
cgcaggcggg tgttcaccag cagcgcgtcg tcctttttga ggatcgccgc caccacgttg 1680  
5 atcccccggt cgagcgctc attgagccac cacacggcgt caaggaaatc gacggcgctc 1740  
tcaatcagta cgatccaccc ctccgcatac tgcgcgcgcg gcagcgctgc cagccgccc 1800  
agggcgatag tgcgtccccc gccaacgccc accccgcccg gcgtctgcgg gttatgaccg 1860  
atcatggctg attcggatg aatgggtctg gtgatggctt ccatcgccac atcgccaatc 1920  
accggcgcgg ctctgtaag atagatgcga gagacatcg tcatcgacca cgggtgtttc 1980  
10 gccagggcct gctccagcgc ggcgaggggc cggcgatat tgtcccgcgt ccctttcatg 2040  
cccgtcgtcg cgacgatccc gctggcaaca aacgcctcgt cctgcgggta gtcggacgcc 2100  
agcgccacct cgggtggtggc gttgccgata tcaatcccgg ctattaacgg catgctgacc 2160  
tccgcttagc ttcctttacg cagcttatgc cgctgctgat acacttccgc cgactcccgg 2220  
acaaaggcgg cattcactgt cgcattgccg gtgtgctcca gctcgtcggc gatcgccagc 2280  
15 agctccgcct gcgaggagcg gaacggggcg agcgcgttat agatagccag aatgcgctcg 2340  
tcaggaatgg cgataagctc cgcgcgcggc cggaaattgc gcgccaccgc atggcgctgc 2400  
atctgctcgg caatctgcgc ctggtactca agggctctggc gggagatccg cacatcctgc 2460  
gggcccacct cgccagagag caccttctcg agggtaatat cggatcaatg tttgccggta 2520  
ggcgtcagga tatgctcggc gcagcgggtg gctaaccgat aatcctgcac gcgcatgggt 2580  
20 ttctcgctca tggctactcc ctactaagt cgatgtgcag ggtgacgggc tcggcgctct 2640  
gcaccacatg tttgggtctt ttgatatgaa atagcgggc tttggccata aatttcggcc 2700  
gcaccatctg atcgttcacc accggcaccg gcgaagggtg ctctttgcgc gcatagcgcg 2760  
cagcgttttt gccaatctgc cggtaggtct ccagcgtcag cagcggcgcc tgggagaaca 2820  
gctccagggt gctgagcggc agcagatcgc gctgatggat gaccgtgggt cccttcgact 2880  
25 ggataccgat gccgatcccc gagccgctca ggttgccgc atcccaggcc ataaaggaga 2940  
cgtcggacgt gcgcagaatg cgcaccaccc gggcgtgaag cccctcttct tccaccccg 3000  
caatcagctc tttgaggatc gcgccatggg gcatatcgat cagagtgtga tgctggtgtt 3060  
tatcgaaggc agggccgacg ccgatcacca ctcatcggc gcgttcatcg gcagaagcta 3120  
ccccgcctc gcgggttttc agggtaaaag agggctgaat ttgggtgtgc tgttgacag 3180  
30 gaataccgcc ttattcaatg gtgtcgggct gaaccacgcc cggaatattt ttgatctccg 3240  
cccagcgttc ggcagagatg cgatagccgg tgcgggccc ctgatagtca ttgatgtcgt 3300  
tgaccgcact caccacctcg aactgccgat cgagaatggc cgaggtctgc aggtaatcgc 3360  
cggtgaccgg ctggcgcagc atattgagaa tattgctggc gatatactca aagccgctgc 3420  
ggctcagcgc gccgacaata tcgaggccgg tgatgttgcg ctcatcatc tcttcaccg 3480  
35 cactcagatc ctccaccacg ttacgcggcg gcatctcgtt gctgccgtgc gcgtagggtg 3540  
cggcctccac ctctcgtcg gcgattggcg gcagccccag ctgcgggaaa accgcctgga 3600  
tcgcccgcgc cgctttctgg cgaatggcaa tggtttccgc ctcggtcacc ggacgcaggc 3660

cgccgtcaac catcaggtca cgctgcagga tgttgaatc atcaaatct tccgcatcga 3720  
 agttcgagcc ggcgaaatg ttgtcgtagt tcggcacgc gctgtagccg gagaaaataa 3780  
 agtcggtgcc cggcagcatc tgcacaggg tgcgcggt gcggcgaata tccgagtggg 3840  
 agaaagtctg gtcgttggtg gacgccactt cgaggtcgag catagaggcg atcaggtttt 3900  
 5 cccgcagcac cccccgaatg cccgacggca cagcgccggt catgccgata cagctcaccg 3960  
 cgccgttttg cagtccctga accccggcgc ctttagtaat gaagatgcag cgcgattcga 4020  
 ggtagagcat cgacttgctc tccgaatagc ccatcagcgc ttcggatccg gtgccggagg 4080  
 tgtagcgc tttcaaccg cgggaggcgt agggcaggc gaggaacgcc tttgaccacg 4140  
 gcgtatcatc gccgtcggta aataccgctt cgggtgccgta gaccgacacc gtctcggcgt 4200  
 10 agctggttaa gccacgcatg cccagctcca gctcgggtggc ctctccacc gagcactgcg 4260  
 tcaacacgcc ggggcgccg cactgcaac cgaccaacag cgccaggcg ttaaacggcg 4320  
 cgtagcgc gataccgacc gtggtctcct gttctgagaa gcccgggatc ccggcctcgg 4380  
 cggcgtcagc ggcaatctgc accgattat ctttgagatt ggtgacgtgg cactggttgg 4440  
 agggggtccg gcgggcagc atcttctgca gcgccatcat catctccacc acgttcatct 4500  
 15 gcgccatcac ctgcaccgct ttggccggcg tgatggcggg agtgatggca atgatctcct 4560  
 cccggctgac gtgaatatcc accagcatac gggctatttc caccgcctcc aggcgcattg 4620  
 cctgctctgt gcgtcaacg ttgatcgcgt aatcgcgat aaatcggtcg atcatgtcaa 4680  
 actggtcccg gcgtttgccg tccagttcga cgatcagacc gttgtccact tttactgaag 4740  
 agaccgggtc aaaggggctg tccatggcga tcagccctc ttcaggccac tcgccaatca 4800  
 20 gccgctctg attgacggg cgctgggcca gtactgcaa tcgttttgat cttttcattg 4860  
 ttcacggct caaaagggtg aatccgcaga cggtagcgaa tacgccggg cagcgtcgtt 4920  
 gccgccggc cattaccggc aatagcggaa ctttaaata ggcagtgggtg aaaaaataa 4980  
 atttaatttc gtttcaattt ggcacacgaa atctaccgac agtttacta tgaaacttta 5040  
 ctccggcggc aaaaaataa aatgtgatcg ccgcgaatga tataaatcaa ttaataaaaa 5100  
 25 acgcccttaa ttacgtttt ccgacgctat ttaacccta tgactaaat catggcgggc 5160  
 gacaaaataa cgctgacaaa aataaagcaa gccaacgaa tggtaatagt tttttactat 5220  
 cgccccctac tgactattcg cgccagcgtt atcctggtgc gggagaga 5268

<210> 10

<211> 607

30 <212> PRT

<213> *Klebsiella pneumoniae*

<400> 10

Met Pro Leu Ile Ala Gly Ile Asp Ile Gly Asn Ala Thr Thr Glu Val

1

5

10

15

Ala Leu Ala Ser Asp Tyr Pro Gln Ala Arg Ala Phe Val Ala Ser Gly  
20 25 30

Ile Val Ala Thr Thr Gly Met Lys Gly Thr Arg Asp Asn Ile Ala Gly  
35 40 45

5 Thr Leu Ala Ala Leu Glu Gln Ala Leu Ala Lys Thr Pro Trp Ser Met  
50 55 60

Ser Asp Val Ser Arg Ile Tyr Leu Asn Glu Ala Ala Pro Val Ile Gly  
65 70 75 80

Asp Val Ala Met Glu Thr Ile Thr Glu Thr Ile Ile Thr Glu Ser Thr  
10 85 90 95

Met Ile Gly His Asn Pro Gln Thr Pro Gly Gly Val Gly Val Gly Val  
100 105 110

Gly Thr Thr Ile Ala Leu Gly Arg Leu Ala Thr Leu Pro Ala Ala Gln  
115 120 125

15 Tyr Ala Glu Gly Trp Ile Val Leu Ile Asp Asp Ala Val Asp Phe Leu  
130 135 140

Asp Ala Val Trp Trp Leu Asn Glu Ala Leu Asp Arg Gly Ile Asn Val  
145 150 155 160

Val Ala Ala Ile Leu Lys Lys Asp Asp Gly Val Leu Val Asn Asn Arg  
20 165 170 175

Leu Arg Lys Thr Leu Pro Val Val Asp Glu Val Thr Leu Leu Glu Gln  
180 185 190

Val Pro Glu Gly Val Met Ala Ala Val Glu Val Ala Ala Pro Gly Gln  
195 200 205

Val Val Arg Ile Leu Ser Asn Pro Tyr Gly Ile Ala Thr Phe Phe Gly  
210 215 220

Leu Ser Pro Glu Glu Thr Gln Ala Ile Val Pro Ile Ala Arg Ala Leu  
225 230 235 240

5 Ile Gly Asn Arg Ser Ala Val Val Leu Lys Thr Pro Gln Gly Asp Val  
245 250 255

Gln Ser Arg Val Ile Pro Ala Gly Asn Leu Tyr Ile Ser Gly Glu Lys  
260 265 270

Arg Arg Gly Glu Ala Asp Val Ala Glu Gly Ala Glu Ala Ile Met Gln  
10 275 280 285

Ala Met Ser Ala Cys Ala Pro Val Arg Asp Ile Arg Gly Glu Pro Gly  
290 295 300

Thr His Ala Gly Gly Met Leu Glu Arg Val Arg Lys Val Met Ala Ser  
305 310 315 320

15 Leu Thr Gly His Glu Met Ser Ala Ile Tyr Ile Gln Asp Leu Leu Ala  
325 330 335

Val Asp Thr Phe Ile Pro Arg Lys Val Gln Gly Gly Met Ala Gly Glu  
340 345 350

Cys Ala Met Glu Asn Ala Val Gly Met Ala Ala Met Val Lys Ala Asp  
20 355 360 365

Arg Leu Gln Met Gln Val Ile Ala Arg Glu Leu Ser Ala Arg Leu Gln  
370 375 380

Thr Glu Val Val Val Gly Gly Val Glu Ala Asn Met Ala Ile Ala Gly  
385 390 395 400

Ala Leu Thr Thr Pro Gly Cys Ala Ala Pro Leu Ala Ile Leu Asp Leu  
405 410 415

Gly Ala Gly Ser Thr Asp Ala Ala Ile Val Asn Ala Glu Gly Gln Ile  
420 425 430

5 Thr Ala Val His Leu Ala Gly Ala Gly Asn Met Val Ser Leu Leu Ile  
435 440 445

Lys Thr Glu Leu Gly Leu Glu Asp Leu Ser Leu Ala Glu Ala Ile Lys  
450 455 460

Lys Tyr Pro Leu Ala Lys Val Glu Ser Leu Phe Ser Ile Arg His Glu  
10 465 470 475 480

Asn Gly Ala Val Glu Phe Phe Arg Glu Ala Leu Ser Pro Ala Val Phe  
485 490 495

Ala Lys Val Val Tyr Ile Lys Glu Gly Glu Leu Val Pro Ile Asp Asn  
500 505 510

15 Ala Ser Pro Leu Glu Lys Ile Arg Leu Val Arg Arg Gln Ala Lys Glu  
515 520 525

Lys Val Phe Val Thr Asn Cys Leu Arg Ala Leu Arg Gln Val Ser Pro  
530 535 540

Gly Gly Ser Ile Arg Asp Ile Ala Phe Val Val Leu Val Gly Gly Ser  
20 545 550 555 560

Ser Leu Asp Phe Glu Ile Pro Gln Leu Ile Thr Glu Ala Leu Ser His  
565 570 575

Tyr Gly Val Val Ala Gly Gln Gly Asn Ile Arg Gly Thr Glu Gly Pro  
580 585 590

Arg Asn Ala Val Ala Thr Gly Leu Leu Leu Ala Gly Gln Ala Asn

595

600

605

&lt;210&gt; 11

&lt;211&gt; 141

5 &lt;212&gt; PRT

<213> *Klebsiella pneumoniae*

&lt;400&gt; 11

Met Ser Glu Lys Thr Met Arg Val Gln Asp Tyr Pro Leu Ala Thr Arg

1

5

10

15

10 Cys Pro Glu His Ile Leu Thr Pro Thr Gly Lys Pro Leu Thr Asp Ile

20

25

30

Thr Leu Glu Lys Val Leu Ser Gly Glu Val Gly Pro Gln Asp Val Arg

35

40

45

Ile Ser Arg Gln Thr Leu Glu Tyr Gln Ala Gln Ile Ala Glu Gln Met

15

50

55

60

Gln Arg His Ala Val Ala Arg Asn Phe Arg Arg Ala Ala Glu Leu Ile

65

70

75

80

Ala Ile Pro Asp Glu Arg Ile Leu Ala Ile Tyr Asn Ala Leu Arg Pro

85

90

95

20 Phe Arg Ser Ser Gln Ala Glu Leu Leu Ala Ile Ala Asp Glu Leu Glu

100

105

110

His Thr Trp His Ala Thr Val Asn Ala Ala Phe Val Arg Glu Ser Ala

115

120

125

Glu Val Tyr Gln Gln Arg His Lys Leu Arg Lys Gly Ser

25

130

135

140

&lt;210&gt; 12

&lt;211&gt; 146

&lt;212&gt; PRT

&lt;213&gt; Klebsiella pneumoniae

5 &lt;400&gt; 12

Met Pro His Gly Ala Ile Leu Lys Glu Leu Ile Ala Gly Val Glu Glu

1 5 10 15

Glu Gly Leu His Ala Arg Val Val Arg Ile Leu Arg Thr Ser Asp Val

20 25 30

10 Ser Phe Met Ala Trp Asp Ala Ala Asn Leu Ser Gly Ser Gly Ile Gly

35 40 45

Ile Gly Ile Gln Ser Lys Gly Thr Thr Val Ile His Gln Arg Asp Leu

50 55 60

Leu Pro Leu Ser Asn Leu Glu Leu Phe Ser Gln Ala Pro Leu Leu Thr

15 65 70 75 80

Leu Glu Thr Tyr Arg Gln Ile Gly Lys Asn Ala Ala Arg Tyr Ala Arg

85 90 95

Lys Glu Ser Pro Ser Pro Val Pro Val Val Asn Asp Gln Met Val Arg

100 105 110

20 Pro Lys Phe Met Ala Lys Ala Ala Leu Phe His Ile Lys Glu Thr Lys

115 120 125

His Val Val Gln Asp Ala Glu Pro Val Thr Leu His Ile Asp Leu Val

130 135 140

Arg Glu

25 145



&lt;210&gt; 13

&lt;211&gt; 555

&lt;212&gt; PRT

&lt;213&gt; Klebsiella pneumoniae

5 &lt;400&gt; 13

Met Lys Arg Ser Lys Arg Phe Ala Val Leu Ala Gln Arg Pro Val Asn

1 5 10 15

Gln Asp Gly Leu Ile Gly Glu Trp Pro Glu Glu Gly Leu Ile Ala Met

20 25 30

10 Asp Ser Pro Phe Asp Pro Val Ser Ser Val Lys Val Asp Asn Gly Leu

35 40 45

Ile Val Glu Leu Asp Gly Lys Arg Arg Asp Gln Phe Asp Met Ile Asp

50 55 60

Arg Phe Ile Ala Asp Tyr Ala Ile Asn Val Glu Arg Thr Glu Gln Ala

15 65 70 75 80

Met Arg Leu Glu Ala Val Glu Ile Ala Arg Met Leu Val Asp Ile His

85 90 95

Val Ser Arg Glu Glu Ile Ile Ala Ile Thr Thr Ala Ile Thr Pro Ala

100 105 110

20 Lys Ala Val Glu Val Met Ala Gln Met Asn Val Val Glu Met Met Met

115 120 125

Ala Leu Gln Lys Met Arg Ala Arg Arg Thr Pro Ser Asn Gln Cys His

130 135 140

Val Thr Asn Leu Lys Asp Asn Pro Val Gln Ile Ala Ala Asp Ala Ala

25 145 150 155 160

Glu Ala Gly Ile Arg Gly Phe Ser Glu Gln Glu Thr Thr Val Gly Ile  
165 170 175

Ala Arg Tyr Ala Pro Phe Asn Ala Leu Ala Leu Leu Val Gly Ser Gln  
180 185 190

5 Cys Gly Arg Pro Gly Val Leu Thr Gln Cys Ser Val Glu Glu Ala Thr  
195 200 205

Glu Leu Glu Leu Gly Met Arg Gly Leu Thr Ser Tyr Ala Glu Thr Val  
210 215 220

Ser Val Tyr Gly Thr Glu Ala Val Phe Thr Asp Gly Asp Asp Thr Pro  
10 225 230 235 240

Trp Ser Lys Ala Phe Leu Ala Ser Ala Tyr Ala Ser Arg Gly Leu Lys  
245 250 255

Met Arg Tyr Thr Ser Gly Thr Gly Ser Glu Ala Leu Met Gly Tyr Ser  
260 265 270

15 Glu Ser Lys Ser Met Leu Tyr Leu Glu Ser Arg Cys Ile Phe Ile Thr  
275 280 285

Lys Gly Ala Gly Val Gln Gly Leu Gln Asn Gly Ala Val Ser Cys Ile  
290 295 300

Gly Met Thr Gly Ala Val Pro Ser Gly Ile Arg Ala Val Leu Ala Glu  
20 305 310 315 320

Asn Leu Ile Ala Ser Met Leu Asp Leu Glu Val Ala Ser Ala Asn Asp  
325 330 335

Gln Thr Phe Ser His Ser Asp Ile Arg Arg Thr Ala Arg Thr Leu Met  
340 345 350

Gln Met Leu Pro Gly Thr Asp Phe Ile Phe Ser Gly Tyr Ser Ala Val  
355 360 365

Pro Asn Tyr Asp Asn Met Phe Ala Gly Ser Asn Phe Asp Ala Glu Asp  
370 375 380

5 Phe Asp Asp Tyr Asn Ile Leu Gln Arg Asp Leu Met Val Asp Gly Gly  
385 390 395 400

Leu Arg Pro Val Thr Glu Ala Glu Thr Ile Ala Ile Arg Gln Lys Ala  
405 410 415

10 Ala Arg Ala Ile Gln Ala Val Phe Arg Glu Leu Gly Leu Pro Pro Ile  
420 425 430

Ala Asp Glu Glu Val Glu Ala Ala Thr Tyr Ala His Gly Ser Asn Glu  
435 440 445

Met Pro Pro Arg Asn Val Val Glu Asp Leu Ser Ala Val Glu Glu Met  
450 455 460

15 Met Lys Arg Asn Ile Thr Gly Leu Asp Ile Val Gly Ala Leu Ser Arg  
465 470 475 480

Ser Gly Phe Glu Asp Ile Ala Ser Asn Ile Leu Asn Met Leu Arg Gln  
485 490 495

20 Arg Val Thr Gly Asp Tyr Leu Gln Thr Ser Ala Ile Leu Asp Arg Gln  
500 505 510

Phe Glu Val Val Ser Ala Val Asn Asp Ile Asn Asp Tyr Gln Gly Pro  
515 520 525

Gly Thr Gly Tyr Arg Ile Ser Ala Glu Arg Trp Ala Glu Ile Lys Asn  
530 535 540

Ile Pro Gly Val Val Gln Pro Asp Thr Ile Glu  
545 550 555

<210> 14

<211> 56

5 <212> DNA

<213> *Escherichia coli*

<400> 14

gctaccatgg cttaaccggt accaaggaga tatcatatgt cagtaccggt tcaaca 56

<210> 15

10 <211> 59

<212> DNA

<213> *Escherichia coli*

<400> 15

gcctcgagtc tagagccgtc gacgggaatt cgagctotta agactgtaaa taaaccacc 59

15 <210> 16

<211> 46

<212> DNA

<213> *Saccharomyces cerevisiae*

<400> 16

20 gcggtaccaa ggaggtatca tatgttcagt agatctacgc tctgct 46

<210> 17

<211> 33

<212> DNA

<213> *Saccharomyces cerevisiae*

&lt;400&gt; 17

gcgaattcga gctcttactc gtccaatttg gcac

34

&lt;210&gt; 18

&lt;211&gt; 35

5 &lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

gcggtaccaa ggaggtatca tatgtcagcc gccgc

35

&lt;210&gt; 19

10 &lt;211&gt; 39

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

gcgaattcga gctcttatga gttcttctga ggcactttg

39

15 &lt;210&gt; 20

&lt;211&gt; 44

&lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;400&gt; 20

20 gcggtaccaa ggaggtatca tatgaccaat aatccccctt cagc

44

&lt;210&gt; 21

&lt;211&gt; 38

&lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;400&gt; 21

gcgaattcga gctcttagaa cagccccaac gggtttatc

38

&lt;210&gt; 22

&lt;211&gt; 20

5 &lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;400&gt; 22

atccccgccgt taaccaccat

20

&lt;210&gt; 23

10 &lt;211&gt; 34

&lt;212&gt; DNA

&lt;213&gt; Escherichia coli

&lt;400&gt; 23

gcggtaccat tggttatccgc tcacaattcc acac

34

15 QBMA0223318

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/23878

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C12P 7/22; C12N 9/02, 9/14, 1/20, 15/00

US CL :435/155, 189, 195, 252.3, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/155, 189, 195, 252.3, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

West; STN files included medline, caplus, uspatfull, embase, scisearch, biosis and biotechds

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|-----------|---|-----------------------|
| A         | TANG et al. Immunochemical Properties of NAD <sup>+</sup> - Linked Glycerol Dehydrogenases from Escherichia coli and Klebsiella pneumoniae. Journal of Bacteriology. December 1982, Vol. 152, No. 3, pages 1169-1174, see the entire article. | 1-12                  |
| A         | BARBIRATO et al. Anaerobic pathways of glycerol dissimilation by Enterobacter agglomerans CNCM 1210: limitations and regulations. Microbiology. 1997, Vol. 143, No. 7, pages 2423-2432.   | 1-12                  |

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

|   |  |
|---|--|
| * Special categories of cited documents:  | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  |
| *A* document defining the general state of the art which is not considered to be of particular relevance  | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone   |
| *B* earlier document published on or after the international filing date  | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *G* document member of the same patent family  |
| *O* document referring to an oral disclosure, use, exhibition or other means  |  |
| *P* document published prior to the international filing date but later than the priority date claimed  |  |

Date of the actual completion of the international search

16 OCTOBER 2000

Date of mailing of the international search report

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

TEKCHAND SAIDHA

Telephone No. (703) 308-0196

**THIS PAGE BLANK (USPTO)**